



Anti-ASAP3 antibody ab187885

2 图像

概述

产品名称	Anti-ASAP3抗体
描述	兔多克隆抗体to ASAP3
宿主	Rabbit
经测试应用	适用于: ICC/IF, IHC-P
种属反应性	与反应: Human
免疫原	Recombinant fragment corresponding to Human ASAP3 aa 650-800. Database link: Q8TDY4
	 Run BLAST with  Run BLAST with
阳性对照	IHC-P: Human smooth muscle tissue; ICC: SiHa whole cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee


Abpromise™承诺保证使用ab187885于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use a concentration of 0.25 - 2 µg/ml. Fixation/Permeabilization: PFA/Triton X-100
IHC-P		1/200 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标	
功能	Promotes cell proliferation.
组织特异性	Highly expressed in primary hepatocarcinoma. Detected in lung, liver and blood leukocytes.
序列相似性	Contains 2 ANK repeats. Contains 1 Arf-GAP domain. Contains 1 PH domain.
细胞定位	Cytoplasm.


图片



Immunohistochemical analysis of paraffin-embedded human smooth muscle tissue labeling ASPA3 in the cytoplasm of smooth muscle cells with ab187885 at a 1/200 dilution.

Performed heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASAP3 antibody (ab187885)



Immunocytochemical analysis of SiHa (Human cervical carcinoma cell line) whole cells labeling ASAP3 in the nucleoplasm and vesicles with ab187885 at 2 µg/ml.

Immunocytochemistry/ Immunofluorescence - Anti-
ASAP3 antibody (ab187885)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors