

Anti-ATF6 antibody [1-7] ab122897



★★★★★ 13 Abreviews 60 References 5 图像

概述

产品名称	Anti-ATF6抗体[1-7]
描述	小鼠单克隆抗体[1-7] to ATF6
宿主	Mouse
特异性	specific to human ATF6 alfa no cross reactivity with mouse ATF6 alfa.
经测试应用	适用于: WB, IP, ICC/IF
种属反应性	与反应: Human 不与反应: Mouse
免疫原	Recombinant fragment (His-tag) corresponding to Human ATF6 (N terminal). Epitope is not determined
阳性对照	293T, HeLa S3 Tet-off, 293 and HeLa cell extracts.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 6 Constituents: 49% PBS, 50% Glycerol
纯化说明	ab122897 was produced from hybridoma cultured in serum-free medium and purified under mild conditions by propriety chromatography processes, then filter sterilized.
克隆	单克隆
克隆编号	1-7

同种型	IgG2a
轻链类型	kappa

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab122897于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (7)	1/500 - 1/1000. Predicted molecular weight: 75 kDa. If clear result not obtained, immunoprecipitation may help.
IP		Use at an assay dependent concentration.
ICC/IF	★★★★★ (4)	1/100.

靶标

功能	Transcription factor that acts during endoplasmic reticulum stress by activating unfolded protein response target genes. Binds DNA on the 5'-CCAC[GA]-3'half of the ER stress response element (ERSE) (5'-CCAAT-N(9)-CCAC[GA]-3') and of ERSE II (5'-ATTGG-N-CCACG-3'). Binding to ERSE requires binding of NF-Y to ERSE. Could also be involved in activation of transcription by the serum response factor.
组织特异性	Ubiquitous.
序列相似性	Belongs to the bZIP family. ATF subfamily. Contains 1 bZIP domain.
结构域	The basic domain functions as a nuclear localization signal. The basic leucine-zipper domain is sufficient for association with the NF-Y trimer and binding to ERSE.
翻译后修饰	During unfolded protein response an approximative 50 kDa fragment containing the cytoplasmic transcription factor domain is released by proteolysis. The cleavage seems to be performed sequentially by site-1 and site-2 proteases. N-glycosylated. The glycosylation status may serve as a sensor for ER homeostasis, resulting in ATF6 activation to trigger the unfolded protein response (UPR). Phosphorylated in vitro by MAPK14/P38MAPK.
细胞定位	Endoplasmic reticulum membrane and Nucleus. Under ER stress the cleaved N-terminal cytoplasmic domain translocates into the nucleus.

图片



Western blot - Anti-ATF6 antibody [1-7] (ab122897)


Lane 1: Wild type HAP1 whole cell lysate (50 µg)

Lane 2: ATF6 knockout HAP1 whole cell lysate (50 µg)

Lanes 1 - 2: Merged signal (red and green). Green - ab122897 observed at 95 kDa. Red - loading control, **ab181602**, observed at 37 kDa.


ab122897 was shown to specifically react with ATF6 in wild-type HAP1 cells along with additional cross-reactive bands. No band was observed when ATF6 knockout samples were used. Wild-type and ATF6 knockout samples were subjected to SDS-PAGE.

Ab122897 and **ab181602** (Rabbit anti GAPDH loading control) were incubated overnight at 4°C at 1/500 dilution and 1/10,000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed (**ab216772**) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed (**ab216777**) secondary antibodies at 1/10,000 dilution for 1hr at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-ATF6 antibody [1-7] (ab122897)

Immunofluorescent analysis of HeLa cells using ab122897 at a dilution of 1/100. The cells were fixed with paraformaldehyde. The secondary antibody was an Alexa Fluor® 488 conjugated goat anti-mouse IgG at a dilution of 1/1000. The antibody staining is shown in the top panel, and the merged image with DAPI counter-staining is shown in the lower panel.



Western blot - Anti-ATF6 antibody [1-7] (ab122897)

All lanes : Anti-ATF6 antibody [1-7] (ab122897) at 1/500 dilution

Lane 1 : 293T cell lysate: control

Lane 2 : 293T cell lysate: treated with 2 µg/ml tunicamycin for 7 hours


Lane 3 : 293T cell lysate: treated with 1mM DTT for 1 hour.

Lane 4 : HeLa S3 Tet-off cell lysate: control

Lane 5 : HeLa S3 Tet-off cell lysate: treated with 1mM DTT for 1 hour.

Predicted band size: 75 kDa

Conversion of precursors pATF6(P) to pATF6(N) in DTT- or tunicamycin-treated cells. ATF6 is constitutively expressed as pATF6a(P) (~90-kDa protein), and converted to pATF6a(N) (>50-kDa protein) in ER-stressed cells.



Western blot - Anti-ATF6 antibody [1-7] (ab122897)

All lanes : Anti-ATF6 antibody [1-7] (ab122897) at 1/500 dilution

Lane 1 : HeLa cell extract: control

Lane 2 : HeLa cell extract: treated with 1mM DTT for 1 hour

Lane 3 : HeLa cell extract: treated with 2 µg/ml tunicamycin for 3 hour

Lane 4 : HeLa cell extract: treated with 2 µg/ml tunicamycin for 7 hour

Lane 5 : 293 cell extract: control


Lane 6 : 293 cell extract: treated with 1mM DTT for 1 hour

Lane 7 : 293 cell extract: treated with 2 µg/ml tunicamycin for 3 hour

Lane 8 : 293 cell extract: treated with 2 µg/ml tunicamycin for 7 hour

Predicted band size: 75 kDa

Conversion of precursors pATF6(P) to pATF6(N) in DTT- or tunicamycin-treated cells. ATF6 is constitutively expressed as pATF6a(P) (~90-kDa protein), and converted to pATF6a(N) (>50-kDa protein) in ER-stressed cells.



Immunoprecipitation - Anti-ATF6 antibody [1-7] (ab122897)

ATF6 was detected by Western blot (Input; lanes 1, 2, 7, and 8)

using **ab122897** (No. 1-7). After immunoprecipitation (IP) with non-related IgG (IP; lanes 3, 4, 9, and 10) or ab122897 (No. 1-7) (IP; lanes 5, 6, 11, and 12), samples were subjected to SDS-PAGE and analyzed by Western blot using **ab122897** (No. 1-7) and anti-mouse IgG antibody (light chain specific).

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors