

### Anti-BarX2 antibody [BAR X2 8A7/1] ab20128

★★★★★ [1 Abreviews](#) [1 References](#)

#### 概述

产品名称	Anti-BarX2抗体[BAR X2 8A7/1]
描述	小鼠单克隆抗体[BAR X2 8A7/1] to BarX2
宿主	Mouse
经测试应用	适用于: ELISA, IHC-P, ICC/IF, WB
种属反应性	与反应: Human
免疫原	Fusion protein: MBP-tagged BarX2 (Human).
常规说明	

BarX2 is a tumour suppressor gene that encodes for a transcription factor known to regulate transcription of cell adhesion molecules in the mouse.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium azide Constituent: 99.98% PBS
纯度	Protein A/G purified
Primary antibody说明	BarX2 is a tumour suppressor gene that encodes for a transcription factor known to regulate transcription of cell adhesion molecules in the mouse.
克隆	单克隆
克隆编号	BAR X2 8A7/1
骨髓瘤	Sp2/0-Ag14

同种型	IgG1
轻链类型	unknown

## 应用

**The Abpromise guarantee** **Abpromise™**承诺保证使用ab20128于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
WB	★★★★★ (1)	Use at an assay dependent concentration.

## 靶标

功能	Transcription factor. Binds optimally to the DNA consensus sequence 5'-YYTAATGRTTTTY-3'. May control the expression of neural adhesion molecules such as L1 or Ng-CAM during embryonic development of both the central and peripheral nervous system. May be involved in controlling adhesive processes in keratinizing epithelia.
组织特异性	Highly expressed in adult salivary gland and at much lower levels in mammary gland, kidney and placenta.
序列相似性	Belongs to the BAR homeobox family. Contains 1 homeobox DNA-binding domain.
细胞定位	Nucleus.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors