abcam

Product datasheet

Biotin Rat IgM, kappa monoclonal [RTK2118] - Isotype control ab35773

概述			
产品名称	生物素大鼠lg M, kappa单 克隆抗体 [RTK2118] -同型对照		
偶联物	Biotin		
经测试应 用	适用于: IHC-Fr, IHC-P, WB, IP, Flow Cyt		
常 规说 明			
	Isotype Control monoclonal antibodies are used to estimate the non-specific binding of target primary antibodies to cell surface antigens. Non-specific binding is due to Fc receptor binding or other protein-protein interactions. Isotype controls show negligible cross-reactivity with cell surface antigens on tissue sections or in cellular preparations. Isotype Controls should be used at identical concentrations and staining conditions as the target primary antibodies.		
	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.		
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As		
性能			
形式	Liquid		
存放 说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.		
存储溶液	pH: 7.20 Preservative: 0.09% Sodium azide Constituent: PBS		
纯 度	Purified IgM		
Isotype control说明	Isotype Control monoclonal antibodies are used to estimate the non-specific binding of target primary antibodies to cell surface antigens. Non-specific binding is due to Fc receptor binding or other protein-protein interactions. Isotype controls show negligible cross-reactivity with cell surface antigens on tissue sections or in cellular preparations. Isotype Controls should be used at identical concentrations and staining conditions as the target primary antibodies.		

克隆	单 克隆
克隆 编号	RTK2118
同种型	lgМ
轻链类型	kappa

应用

The Abpromise guarantee Abpromise™承诺保证使用ab35773于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说 明
IHC-Fr		Use at an assay dependent dilution.
IHC-P		Use at an assay dependent dilution.
WB		Use at an assay dependent dilution.
IP		Use at an assay dependent dilution.
Flow Cyt		Use at an assay dependent dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors