# abcam

### Product datasheet

## Anti-C1orf123 antibody ab122865

1 References 3 图像

概述

产**品名称** Anti-C1orf123抗体

描述 兔多克隆抗体to C1orf123

宿主 Rabbit

经测试应用 适用于: IHC-P, WB, ICC/IF

种属反应性 与反应: Human

免疫原 antigen sequence: IALQLKATLE NITNLRPVGE DFRWYLKMKC GNCGEISDKW QYIRLMDSVA

LKGGRGSASM VQKCKLCARE, corresponding to amino acids 4-73 of Human C1orf123.

(Q9NWV4)

Run BLAST with EXPASY Run BLAST with S NCBI

阳性对照 Human liver tissue.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

**存放说明** Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

**存储溶液** pH: 7.20

Preservative: 0.02% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)

纯**度** Immunogen affinity purified

**克隆** 多克隆

**同种型** lgG

应用

1

#### The Abpromise guarantee

#### Abpromise™承诺保证使用ab122865于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
IHC-P		1/20 - 1/50. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 µg/ml.
ICC/IF		Use a concentration of 0.25 - 2 µg/ml.

#### 靶标


#### 序列相似性

Belongs to the UPF0587 family.

#### 图片




Immunocytochemistry/ Immunofluorescence - Anti-C1orf123 antibody (ab122865) Immunofluorescent staining of Human cell line U-2 OS shows positivity in cytoplasm. Recommended concentration of ab122865 1-4 µg/ml. Cells treated with PFA/Triton X-100.



Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10

Lane 2: Negative control (vector only transfected HEK293T lysate)

Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-C1orf123 antibody (ab122865)

ab122865 at 1/20 dilution staining C1orf123 in paraffin-embedded Human liver tissue by Immunohistochemistry.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors