


Anti-Caspase-3 antibody ab13847

敲除验证

★★★★☆ [56 Abreviews](#) [1118 References](#) [2 图像](#)

概述

产品名称	Anti-Caspase-3抗体
描述	兔多克隆抗体to Caspase-3
宿主	Rabbit
特异性	<p>Stimulation is required to allow detection of the 17kDa cleaved form of the protein. Please see images below for recommended treatment conditions and positive controls.</p> <p>ab13847 recognizes a cleaved form of Caspase 3 (~17 kDa) after apoptosis has been induced in wildtype cells and not Caspase 3 knockout cells.</p> <p>Some customers have used this antibody successfully in IHC-P however our latest tests were unsuccessful and therefore we can no longer guarantee this application. We would recommend ab32351 and ab184787 as an alternative product for this application.</p>

经测试应用	适用于: WB
种属反应性	<p>与反应: Human</p> <p>预测可用于: Mouse, Rat, Dog, Pig, Xenopus laevis, Drosophila melanogaster, Indian muntjac, Zebrafish, Rhesus monkey, Chinese hamster, Common marmoset, Schmidtea mediterranea, Salvelinus alpinus </p>

免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Hap-1 WT, Human Caspase 3 (active) Recombinant Protein ICC: HeLa cells treated with 1 mM staurosporine

常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>
------	---

性能

形式	Liquid
----	--------

存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab13847于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (15)	1/500. Detects a band of approximately 17, 34 kDa (predicted molecular weight: 17, 34 kDa).

靶标

功能	Involved in the activation cascade of caspases responsible for apoptosis execution. At the onset of apoptosis it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7 and -9. Involved in the cleavage of huntingtin.
组织特异性	Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.
序列相似性	Belongs to the peptidase C14A family.
翻译后修饰	Cleavage by granzyme B, caspase-6, caspase-8 and caspase-10 generates the two active subunits. Additional processing of the propeptides is likely due to the autocatalytic activity of the activated protease. Active heterodimers between the small subunit of caspase-7 protease and the large subunit of caspase-3 also occur and vice versa. S-nitrosylated on its catalytic site cysteine in unstimulated human cell lines and denitrosylated upon activation of the Fas apoptotic pathway, associated with an increase in intracellular caspase activity. Fas therefore activates caspase-3 not only by inducing the cleavage of the caspase zymogen to its active subunits, but also by stimulating the denitrosylation of its active site thiol.
细胞定位	Cytoplasm.

图片



Western blot - Anti-Caspase-3 antibody (ab13847)

Lane 1: Wild-type HAP1 cell lysate + Staurosporine ([ab146588](#)) (1 μ M for 4h)


Lane 2: Wild-type HAP1 cell lysate

Lane 3: Caspase-3 knockout HAP1 cell lysate + Staurosporine ([ab146588](#)) (1 μ M for 4h)

Lane 4: Caspase-3 knockout HAP1 cell lysate

Lanes 1 - 4: Merged signal (red and green). Green - ab13847 observed at 17 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab13847 was shown to recognise Caspase 3 when Caspase 3 knockout samples were used, along with additional cross-reactive bands. Wild-type and Caspase 3 knockout samples (\pm staurosporine treatment) were subjected to SDS-PAGE. ab13847 and [ab8245](#) (loading control to GAPDH) were diluted to 1/500 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with [Goat anti-Rabbit IgG H&L \(IRDye® 800CW\) preadsorbed \(ab216773\)](#) and [Goat anti-Mouse IgG H&L \(IRDye® 680RD\) preadsorbed \(ab216776\)](#) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Caspase-3 antibody (ab13847)

All lanes : Anti-Caspase-3 antibody (ab13847) at 1 μ g/ml

Lane 1 : Human Caspase 3 (active) Recombinant Protein

Lane 2 : Human Pro Caspase 3 (inactive) Recombinant Protein

Lysates/proteins at 0.1 μ g per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 17, 34 kDa

Observed band size: 17, 32 kDa

Exposure time: 8 minutes

Caspase 3 exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce large (17kDa) and small (12kDa) subunits. These subunits dimerize to form the active enzyme. ab13847 specifically detects the large active subunit (17kDa) and the inactive pro Caspase 3 (32 kDa).

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Bovine Serum Albumin before being incubated with ab13847 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution.

Secondary antibody - **Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody**

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors