

### Anti-CaSR antibody [5C10, ADD] ab19347

★★★★★ [5 Abreviews](#) [56 References](#) [5 图像](#)

#### 概述

产品名称	Anti-CaSR抗体[5C10, ADD]
描述	小鼠单克隆抗体[5C10, ADD] to CaSR
宿主	Mouse
经测试应用	适用于: IHC-P, Flow Cyt, IHC-Fr
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide within Human CaSR aa 200-300. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please <b><a href="#">contact</a></b> our Scientific Support team to discuss your requirements.

 [Run BLAST with](#)

 [Run BLAST with](#)

#### 常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: 0.05% Sodium azide Constituents: PBS, 0.1% BSA
纯度	Immunogen affinity purified
克隆	单克隆
克隆编号	5C10, ADD
同种型	IgG2a

## 应用


**The Abpromise guarantee**      **Abpromise™**承诺保证使用ab19347于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (1)	Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Flow Cyt		Use 1µg for 10 <sup>6</sup> cells. <b>ab170191</b> - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
IHC-Fr	★★★★★ (3)	Use a concentration of 2 µg/ml.

## 靶标

功能	Senses changes in the extracellular concentration of calcium ions. The activity of this receptor is mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system.
组织特异性	Expressed in the temporal lobe, frontal lobe, parietal lobe, hippocampus, and cerebellum. Also found in kidney, lung, liver, heart, skeletal muscle, placenta.
疾病相关	<p>Defects in CASR are the cause of familial hypocalciuric hypercalcemia type 1 (FHH) [MIM:145980]. FHH is characterized by altered calcium homeostasis. Affected individuals exhibit mild or modest hypercalcemia, relative hypocalciuria, and inappropriately normal PTH levels.</p> <p>Defects in CASR are the cause of neonatal severe primary hyperparathyroidism (NSHPT) [MIM:239200]. NSHPT is a rare autosomal recessive life-threatening disorder characterized by very high serum calcium concentrations, skeletal demineralization, and parathyroid hyperplasia. In some instances NSHPT has been demonstrated to be the homozygous form of FHH.</p> <p>Defects in CASR are a cause of familial isolated hypoparathyroidism (FIH) [MIM:146200]; also called autosomal dominant hypoparathyroidism or autosomal dominant hypocalcemia. FIH is characterized by hypocalcemia and hyperphosphatemia due to inadequate secretion of parathyroid hormone. Symptoms are seizures, tetany and cramps. An autosomal recessive form of FIH also exists.</p> <p>Defects in CASR are the cause of idiopathic generalized epilepsy type 8 (IGE8) [MIM:612899]; also known as EIG8. A disorder characterized by recurring generalized seizures in the absence of detectable brain lesions and/or metabolic abnormalities. Seizure types are variable, but include myoclonic seizures, absence seizures, febrile seizures, complex partial seizures, and generalized tonic-clonic seizures.</p> <p>Note=Homozygous defects in CASR can be a cause of primary hyperparathyroidism in adulthood. Patients suffer from osteoporosis and renal calculi, have marked hypercalcemia and increased serum PTH concentrations.</p>
序列相似性	Belongs to the G-protein coupled receptor 3 family.
翻译后修饰	N-glycosylated. Ubiquitinated by RNF19A; which induces proteasomal degradation.
细胞定位	Cell membrane.




Immunohistochemistry (Frozen sections) - Anti-CaSR antibody [5C10, ADD] (ab19347)

Immunohistochemical analysis of 4% formalin fixed frozen mouse stomach tissue labeling CaSR with ab19347 at 1/100 dilution overnight at 4°C, followed by fluorophore-conjugated goat anti-mouse IgG secondary antibody for 2 hours at RT.


Fresh stomach tissue was fixed in 4% formalin for 1 hour and then incubated overnight at in 25% sucrose before embedding in tissue freezing medium. Antigen retrieval was carried out on 8µm cryo-sections by incubating in sodium citrate buffer for 45 minutes at 4°C, immersing in sodium citrate buffer for 10 minutes at 100°C before washing 3 times for 5 minutes each in 1X PBS. Sections were then blocked in blocking buffer (0.3% Triton X-100 in 1X PBS containing 10% normal goat serum) for 30 minutes at RT before staining with ab19347.

Sections were also stained with DAPI nuclear stain (blue). Positive cells (green) were found at the base of the antral glands in the mouse stomach.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CaSR antibody [5C10, ADD] (ab19347)

Immunohistochemistry was performed on normal biopsies of deparaffinized Human kidney tissue. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a mouse monoclonal antibody recognizing CaSR ab19347 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.




Flow Cytometry - Anti-CaSR antibody [5C10, ADD]  
(ab19347)

Overlay histogram showing SH-SY5Y cells stained with ab19347 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab19347, 1 µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-mouse IgG (H+L) (**ab150113**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2a [ICIGG2A] (**ab91361**, 1 µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive result in 80% methanol (5 min) fixed SH-SY5Y cells used under the same conditions. Please note that Abcam do not have any data for use of this antibody on non-fixed cells. We welcome any customer feedback.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CaSR antibody [5C10, ADD] (ab19347)

Immunohistochemistry was performed on normal biopsies of deparaffinized Human brain tissue. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a mouse monoclonal antibody recognizing CaSR ab19347 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.



Immunohistochemistry (Frozen sections) - Anti-CaSR antibody [5C10, ADD] (ab19347)

This image is courtesy of an anonymous Abreview

ab19347 at 1/100 staining rat brain (cerebral cortex) tissue sections by IHC-Fr. The tissue was paraformaldehyde fixed and blocked with serum before incubation with the primary antibody for 24 hours at 4°C. A biotinylated horse anti-mouse IgG was used as the secondary.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors