

# Anti-CD45 antibody [OX30] ab6329

**2 References**   **1 图像**

### 概述

产品名称	Anti-CD45抗体[OX30]
描述	小鼠单克隆抗体[OX30] to CD45
宿主	Mouse
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Rat
免疫原	<p><b>This product was produced with the following immunogens:</b></p> <p>Full length protein. This information is proprietary to Abcam and/or its suppliers.</p> <p>Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.</p>
阳性对照	Flow Cyt: Lewis rat splenocytes.
常规说明	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium azide Constituent: PBS
纯度	Protein G purified
纯化说明	Purified from TCS.
克隆	单克隆

克隆编号	OX30
同种型	IgG2a
轻链类型	kappa

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab6329于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use a concentration of 0.2 µg/ml.

靶标

功能	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN.
疾病相关	Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+)) SCID [MIM:608971]. A form of severe combined immunodeficiency (SCID), a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients present in infancy recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development. Genetic variations in PTPRC are involved in multiple sclerosis susceptibility (MS) [MIM:126200]. MS is a neurodegenerative disorder characterized by the gradual accumulation of focal plaques of demyelination particularly in the periventricular areas of the brain. Peripheral nerves are not affected. Onset usually in third or fourth decade with intermittent progression over an extended period. The cause is still uncertain.
序列相似性	Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. Contains 2 fibronectin type-III domains. Contains 2 tyrosine-protein phosphatase domains.
结构域	The first PTPase domain interacts with SKAP1.
翻译后修饰	Heavily N- and O-glycosylated.
细胞定位	Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts.

图片



Lewis rat splenocytes stained with ab6329 (right) or mouse IgG2ak (left). Lewis rat splenocytes were incubated for 30 min on ice in 10% rat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab6329) or mouse IgG2ak Isotype (**ab18450**) ( $1 \times 10^6$  in 100  $\mu$ l at 0.2  $\mu$ g/ml) for 30 min on ice.

The secondary antibody Goat Anti-Mouse IgG H&L (Alexa Fluor<sup>®</sup> 488, pre-adsorbed) (**ab150117**) was used at 1/2000 dilution for 30 min at 4°C. The cells were simultaneously stained with CD3 APC antibody.

Acquisition of >30,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter. Events were gated on viable lymphocytes.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors