abcam

Product datasheet

Anti-CD62L antibody [EPR22188-297] ab218829



重组 RabMAb

2 图像

概述

产品名称 Anti-CD62L抗体[EPR22188-297]

描述 兔单克隆抗体[EPR22188-297] to CD62L

宿主 Rabbit

经测试应用 适用于: Flow Cyt

不适用于: ICC/IF,IHC-P or WB

种属反应性 **与反应:** Rat

免疫原 Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.

阳性对照 Flow Cyt: Rat primary splenocytes.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

存储溶液 pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

纯度 Protein A purified

克隆 单克降

克隆编号 EPR22188-297

同种型 ΙgG

The Abpromise guarantee

Abpromise™承诺保证使用ab218829于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		1/500.

应用说明

Is unsuitable for ICC/IF,IHC-P or WB.

靶标


相关性

CD62L is a cell surface component that is a member of a family of adhesion/homing receptors. These play important roles in leukocyte-endothelial cell interactions. CD62L is composed of multiple domains: one homologous to lectins, one to epidermal growth factor, and two to the consensus repeat units found in C3/C4 binding proteins.

细胞定位

Cell Membrane; single-pass type I membrane protein.

图片




Flow Cytometry - Anti-CD62L antibody [EPR22188-297] (ab218829)

Flow cytometric analysis of rat primary splenocytes labeling CD62L with ab218829 at 1/500 dilution (top right panel), compared with a Rabbit lgG, monoclonal [EPR25A] - Isotype Control (ab172730) (Upper left). Goat Anti-Rabbit lgG H&L (Alexa Fluor[®] 488) (ab150077) at 1/2000 dilution was used as the secondary antibody.

Top panel: Cells were stained with Alexa Fluor[®] 647-conjugated CD3 and rabbit lgG (Left) or anti-CD62L RabMab (Right). Data shown was gated on viable cells.

Down regulation of CD62L on activated T cells (CD3+FSC^{high}, blue) compared to Naïve T cells (CD3+CD62L+, red) was shown on the bottom panel.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors