

Anti-CD62P antibody [EPR2146(2)] ab182135

重组 RabMAb

2 References **6 图像**

概述

产品名称	Anti-CD62P抗体[EPR2146(2)]
描述	兔单克隆抗体[EPR2146(2)] to CD62P
宿主	Rabbit
经测试应用	适用于: IHC-P, WB, IP 不适用于: Flow Cyt or ICC/IF
种属反应性	与反应: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Human fetal liver, Human placenta and plasma lysates; Human spleen and gliocytoma tissues.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2146(2)
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab182135于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
IHC-P		1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		1/1000 - 1/10000. Detects a band of approximately 91 kDa (predicted molecular weight: 91 kDa).
IP		1/30.

应用说明 Is unsuitable for Flow Cyt or ICC/IF.

靶标

功能	Ca(2+)-dependent receptor for myeloid cells that binds to carbohydrates on neutrophils and monocytes. Mediates the interaction of activated endothelial cells or platelets with leukocytes. The ligand recognized is sialyl-Lewis X. Mediates rapid rolling of leukocyte rolling over vascular surfaces during the initial steps in inflammation through interaction with PSGL1.
组织特异性	Stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. Upon cell activation by agonists, P-selectin is transported rapidly to the cell surface.
疾病相关	Defects in SELP may be a cause of susceptibility to ischemic stroke (ISCHSTR) [MIM:601367]; also known as cerebrovascular accident or cerebral infarction. A stroke is an acute neurologic event leading to death of neural tissue of the brain and resulting in loss of motor, sensory and/or cognitive function. Ischemic strokes, resulting from vascular occlusion, is considered to be a highly complex disease consisting of a group of heterogeneous disorders with multiple genetic and environmental risk factors.
序列相似性	Belongs to the selectin/LECAM family. Contains 1 C-type lectin domain. Contains 1 EGF-like domain. Contains 9 Sushi (CCP/SCR) domains.
细胞定位	Membrane.


图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD62P antibody [EPR2146(2)] (ab182135)

Immunohistochemical analysis of paraffin-embedded Human gliocytoma tissue labeling CD62P with ab182135 at 1/500 dilution followed by pre-diluted HRP-conjugated secondary antibody and counter-stained with Hematoxylin.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD62P antibody [EPR2146(2)] (ab182135)


Immunohistochemical analysis of paraffin-embedded Human spleen tissue labeling CD62P with ab182135 at 1/500 dilution followed by pre-diluted HRP-conjugated secondary antibody and counter-stained with Hematoxylin.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot analysis of immunoprecipitation pellet from Human plasma immunoprecipitated using ab182135 at 1/30 dilution. Secondary antibody Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000.

Immunoprecipitation - Anti-CD62P antibody [EPR2146(2)] (ab182135)




Anti-CD62P antibody [EPR2146(2)] (ab182135) at 1/10000 dilution + Human plasma lysate at 20 µg

Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 91 kDa

Western blot - Anti-CD62P antibody [EPR2146(2)] (ab182135)



Western blot - Anti-CD62P antibody [EPR2146(2)] (ab182135)

All lanes : Anti-CD62P antibody [EPR2146(2)] (ab182135) at 1/2000 dilution

Lane 1 : Human fetal liver lysate

Lane 2 : Human placenta lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), HRP at 1/1000 dilution

Predicted band size: 91 kDa

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-CD62P antibody [EPR2146(2)] (ab182135)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors