abcam

Product datasheet

Anti-CD81 antibody [1D6] - Low endotoxin, Azide free ab35026

★★★★★ 1 Abreviews 5 References 1 图像

概述

产品名称	Anti-CD81 抗体 [1D6] - Low endotoxin, Azide free	
描述	小鼠单克隆抗体[1D6] to CD81 - Low endotoxin, Azide free	
宿主	Mouse	
特异性	ab35026 recognises CD81, a 26kD cell surface antigen and a member of the tetraspanin family.	
经 测 试应 用	适用于: Flow Cyt 不适用于: WB	
种属反 应性	与反应: Human	
免疫原	Tissue, cells or virus corresponding to Human CD81. Aggregated OCHLY8 cells.	
阳性 对照	Flow Cyt: HAP1-WT cells.	
常 规说 明	ab35026 is a potent CD81 reagent, induces homotypic adhesion and has powerful anti proliferative effects.	
	Endotoxin Level: <0.01 Eu/ug (low endotoxin grade).	
	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.	
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As	

性能	
形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.40 Constituent: PBS
无载体	是一一是一个人们的问题,我们们的问题,我们们的问题,我们们的问题,我们们就能能能能能。
纯 度	Protein G purified

纯化说明	This antibody was purified from tissue culture supernatant.	
Primary antibody说明	Ab35026 is a potent CD81 reagent, induces homotypic adhesion and has powerful anti proliferative effects. Endotoxin Level: <0.01 Eu/ug (low endotoxin grade).	
克隆	单 克隆	
克 隆 编号	1D6	
骨髓瘤	P3x63-Ag8.653	
同种型	lgG1	


应用

The Abpromise guarantee Abpromise™承诺保证使用ab35026于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明	
Flow Cyt		1/5 - 1/10. Use 10μ of the suggested working dilution to label 10^6 cells in 100μ l. ab170190 - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.	
应用说明	ls unsuitable for WB.		
靶标			
功能	May play an important role in the regulation of lymphoma cell growth. Interacts with a 16-kDa Leu- 13 protein to form a complex possibly involved in signal transduction. May acts a the viral receptor for HCV.		
组织 特异性	Hematolymphoid, neuroectodermal and mesenchymal tumor cell lines.		
疾病相关	Defects in CD81 are the cause of immunodeficiency common variable type 6 (CVID6) [MIM:613496]; also called antibody deficiency due to CD81 defect. CVID6 is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antigen. The defect results from a failure of B-cell differentiation and impaired secretion of immunoglobulins; the numbers of circulating B cells is usually in the normal range, but can be low.		
序列相似性	Belongs to the tetraspanin (TM4SF) family.		
翻 译 后修 饰	Not glycosylated.		
细 胞定位	Membrane.		

图片



Flow Cytometry - Anti-CD81 antibody [1D6] - Low endotoxin, Azide free (ab35026)

Overlay histogram showing HAP1 wildtype (green line) and HAP1-CD81 knockout cells (red line) stained with ab35026. The cells were fixed with 4% formaldehyde (10 min) and then permeabilized with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab35026, 0.1µg/ml) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-mouse IgG (H&L) presorbed (**ab150117**) at 1/2000 dilution for 30 min at 22°C.

A mouse IgG1 isotype control antibody (**ab170190**) was used at the same concentration and conditions as the primary antibody (HAP1 wildtype - black line, HAP1-CD81 knockout - grey line). Unlabelled sample was also used as a control (this line is not shown for the purpose of simplicity).

Acquisition of >5,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter.

This antibody can also be used in HAP1 cells fixed with 80% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 15 min under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors