abcam

Product datasheet

Anti-CIP75 antibody ab106443

★★★★★ 2 Abreviews 4 References 2 图像

概述

产**品名称** Anti-CIP75抗体

描述 兔多克隆抗体to CIP75

宿主 Rabbit

经测试应用 适用于: IHC-P, WB

种属反应性 与反应: Mouse

免疫原 Synthetic peptide corresponding to Human CIP75 (N terminal).

Database link: Q9NRR5

阳性对照 3T3 cell lysate, Mouse kidney tissue.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

常规说明

形式 Liquid

存放说明 Shipped at 4°C. Store at 4°C (stable for up to 12 months).

存储溶液 pH: 7.2

Preservative: 0.02% Sodium azide

Constituent: PBS

纯**度** Immunogen affinity purified

克隆 多克隆 **同种型** IqG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab106443于以下的经测试应用

ALTERNATION OF A STATE OF A STATE

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use a concentration of 5 μ g/ml. Perform heat mediated antigen retrieval via the microwave method before commencing with IHC staining protocol.
WB	*****(1)	Use a concentration of 1 - 2 μg/ml. Predicted molecular weight: 64 kDa.

靶标

组织**特异性** Highly expressed in pancreas, kidney, skeletal muscle, heart and throughout the brain, and at

lower levels in placenta, lung and liver.


序列相似性 Contains 1 UBA domain.

Contains 1 ubiquitin-like domain.

翻译后修饰 Phosphorylated upon DNA damage, probably by ATM or ATR.

细胞定位 Nucleus. Cytoplasm. Endoplasmic reticulum. May associate with the endoplasmic reticulum.

图片




Western blot - Anti-CIP75 antibody (ab106443)

Lane 1: Anti-CIP75 antibody (ab106443) at 1 μg/ml **Lane 2**: Anti-CIP75 antibody (ab106443) at 2 μg/ml

All lanes: 3T3 cell lysate

Lysates/proteins at 15 µg per lane.

Predicted band size: 64 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CIP75 antibody (ab106443)

ab106443, at 5 ug/ml, staining CIP75 in Mouse kidney tissue by immunohistochemistry.

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors