

Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867] ab133319

敲除验证 重组 RabMAb


★★★★★ 1 Abreviews 8 References 5 图像

概述

产品名称	Anti-COX1 / Cyclooxygenase 1抗体[EPR5867]
描述	兔单克隆抗体[EPR5867] to COX1 / Cyclooxygenase 1
宿主	Rabbit
经测试应用	适用于: WB, IHC-P 不适用于: Flow Cyt, ICC/IF or IP
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Human COX1/ Cyclooxygenase 1 aa 500-600. The exact sequence is proprietary.
阳性对照	C2C12, A431, NIH 3T3, Neuro 2a, and L6 cell lysates; Human stomach tissue.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information <a href="#">see here</a> .  Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
解离常数 (K <sub>D</sub> )	K <sub>D</sub> = 8.00 x 10 <sup>-11</sup> M



存储溶液	pH: 7.2 Preservative: 0.05% Sodium azide
------	---

	Constituents: 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR5867
同种型	IgG

应用

**The Abpromise guarantee**      **Abpromise™**承诺保证使用ab133319于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB	★★★★☆ (1)	1/1000 - 1/10000. Detects a band of approximately 65-70 kDa (predicted molecular weight: 68 kDa).
IHC-P		1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

**应用说明**      Is unsuitable for Flow Cyt, ICC/IF or IP.

靶标

功能	May play an important role in regulating or promoting cell proliferation in some normal and neoplastically transformed cells.
通路	Lipid metabolism; prostaglandin biosynthesis.
序列相似性	Belongs to the prostaglandin G/H synthase family. Contains 1 EGF-like domain.
细胞定位	Microsome membrane. Endoplasmic reticulum membrane.

图片



**All lanes :** Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867] (ab133319) at 1/1000 dilution

**Lane 1 :** Wild-type A431 cell lysate

**Lane 2 :** PTGS1 knockout A431 cell lysate

**Lane 3 :** C2C12 cell lysate

**Lane 4 :** MOLT-4 cell lysate

Lysates/proteins at 20 µg per lane.

## Secondary

**All lanes :** Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution


Performed under reducing conditions.

**Predicted band size:** 68 kDa

**Observed band size:** 70 kDa

False colour image of Western blot: Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red.


In Western blot, ab133319 was shown to bind specifically to COX1 / Cyclooxygenase 1. A band was observed at 70 kDa in wild-type A431 cell lysates with no signal observed at this size in PTGS1 knockout cell line [ab270477](#) (knockout cell lysate [ab270500](#)). To generate this image, wild-type and PTGS1 knockout A431 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867] (ab133319)

Immunohistochemical analysis of Paraffin Embedded Human stomach tissue labelling COX1/Cyclooxygenase 1 with ab133319 at 1/50 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867] (ab133319)

**All lanes :** Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867] (ab133319) at 1/1000 dilution

**Lane 1 :** C2C12 cell lysate

**Lane 2 :** A431 cell lysate

**Lane 3 :** NIH/3T3 cell lysate

**Lane 4 :** Neuro 2a cell lysate


**Lane 5 :** L6 cell lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit HRP at 1/2000 dilution

**Predicted band size:** 68 kDa



OI-RD Scanning - Anti-COX1 / Cyclooxygenase 1  
 antibody [EPR5867] (ab133319)

Equilibrium dissociation constant ( $K_D$ )

Learn more about  $K_D$

[Click here to learn more about  \$K\_D\$](#)

### Why choose a recombinant antibody?



**Research with confidence**  
 Consistent and reproducible results



**Long-term and scalable supply**  
 Recombinant technology



**Success from the first experiment**  
 Confirmed specificity



**Ethical standards compliant**  
 Animal-free production

Anti-COX1 / Cyclooxygenase 1 antibody [EPR5867]  
 (ab133319)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors