




Anti-CREB + ICER antibody ab5803

★★★★★ [3 Abreviews](#) [11 References](#) [7 图像](#)

概述

产品名称	Anti-CREB + ICER抗体
描述	兔多克隆抗体to CREB + ICER
宿主	Rabbit
特异性	ab5803 detects both the phosphorylated and non-phosphorylated forms of cyclic-AMP response element binding protein (CREB) from rat cells.
经测试应用	适用于: WB, ICC/IF, IHC-P
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Cow, Dog, Zebrafish 
免疫原	Synthetic peptide corresponding to Human CREB aa 123-136. Sequence: KRREILSRPSYRK (Peptide available as ab5860) <div>  Run BLAST with  Run BLAST with </div>
阳性对照	WB: GH4 cell extract. ICC/IF: SK-N-MC cells, Neuro-2a cells. IHC-P: Mouse brain tissue, Human glioma, Human lung adenocarcinoma.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 99% PBS

纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™承诺保证使用ab5803于以下的经测试应用**

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★☆ (2)	Use a concentration of 2 µg/ml. Detects a band of approximately 43 kDa (predicted molecular weight: 43 kDa).Can be blocked with CREB + ICER peptide (ab5860) .
ICC/IF	★★★★☆ (1)	1/50 - 1/500.
IHC-P		1/20 - 1/200.


靶标

细胞定位 CREB: Nucleus. CREM: Nucleus.

图片




Shows a Western blot of CREB on GH4 cell extract using ab5803.




Immunofluorescent analysis of CREB (green) showing staining in the nucleus of SK-N-MC cells (right) compared to a negative control without primary antibody (left). Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with a CREB polyclonal antibody (ab5803) in 3% BSA-PBS at a dilution of 1:200 and incubated overnight at 4°C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody in PBS at room temperature in the dark. Actin was stained using Alexa Fluor 554

(red) and nuclei were stained with Hoechst or DAPI (blue). Images were taken at a magnification of 60x.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CREB + ICER antibody (ab5803)

Immunohistochemistry analysis of CREB showing staining in the nucleus of paraffin-embedded mouse brain tissue (right) compared to a negative control without primary antibody (left). To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 min. Following antigen retrieval, tissues were blocked in 3% H₂O₂-methanol for 15 min at room temperature, washed with ddH₂O and PBS, and then probed with a CREB polyclonal antibody (ab5803) diluted in 3% BSA-PBS at a dilution of 1:50 overnight at 4°C in a humidified chamber. Tissues were washed extensively in PBST and detection was performed using an HRP-conjugated secondary antibody followed by colorimetric detection using a DAB kit. Tissues were counterstained with hematoxylin and dehydrated with ethanol and xylene to prep for mounting.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CREB + ICER antibody (ab5803)

Immunohistochemistry analysis of CREB showing staining in the nucleus of paraffin-embedded Human glioma (right) compared to a negative control without primary antibody (left). To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 min. Following antigen retrieval, tissues were blocked in 3% H₂O₂-methanol for 15 min at room temperature, washed with ddH₂O and PBS, and then probed with a CREB polyclonal antibody (ab5803) diluted in 3% BSA-PBS at a dilution of 1:100 overnight at 4°C in a humidified chamber. Tissues were washed extensively in PBST and detection was performed using an HRP-conjugated secondary antibody followed by colorimetric detection using a DAB kit. Tissues were counterstained with hematoxylin and dehydrated with ethanol and xylene to prep for mounting.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CREB + ICER antibody (ab5803)

Immunohistochemistry analysis of CREB showing staining in the nucleus of paraffin-embedded Human lung adenocarcinoma (right) compared to a negative control without primary antibody (left). To expose target proteins, antigen retrieval was performed using 10mM sodium citrate (pH 6.0), microwaved for 8-15 min. Following antigen retrieval, tissues were blocked in 3% H₂O₂-methanol for 15 min at room temperature, washed with ddH₂O and PBS, and then probed with a CREB polyclonal antibody (ab5803) diluted in 3% BSA-PBS at a dilution of 1:100 overnight at 4°C in a humidified chamber. Tissues were washed extensively in PBST and detection was performed using an HRP-conjugated secondary antibody followed by colorimetric detection using a DAB kit. Tissues were counterstained with hematoxylin and dehydrated with ethanol and xylene to prep for mounting.



Immunocytochemistry/ Immunofluorescence - Anti-CREB + ICER antibody (ab5803)

Immunofluorescent analysis of CREB (green) showing staining in the nucleus of Neuro-2a cells (right) compared to a negative control without primary antibody (left). Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with a CREB polyclonal antibody (ab5803) in 3% BSA-PBS at a dilution of 1:100 and incubated overnight at 4°C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody in PBS at room temperature in the dark. Nuclei were stained with Hoechst or DAPI (blue). Images were taken at a magnification of 60x.



Western blot - Anti-CREB + ICER antibody (ab5803)

This image is courtesy of Richard D'Mello, Kings College, London

All lanes : Anti-CREB + ICER antibody (ab5803) at 1/500 dilution

All lanes : hippocampal lysate

Lysates/proteins at 40 µg per lane.

Secondary

All lanes : Donkey Anti-Rabbit IR800-linked conjugated to IRDye 800CW at 1/15000 dilution

Performed under reducing conditions.

Predicted band size: 43 kDa

Observed band size: 43 kDa

Additional bands at: 55 kDa (possible non-specific binding)

Exposure time: 5 minutes

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors