


Anti-CREB antibody [LB9] ab178322

敲除 验证

★★★★☆ **1 Abreviews** **15 References** **8 图像**

概述

产品名称	Anti-CREB抗体[LB9]
描述	小鼠单克隆抗体[LB9] to CREB
宿主	Mouse
经测试应用	适用于: ICC/IF, WB, IP, IHC-P
种属反应性	与反应: Mouse, Rat, Human, African green monkey 预测可用于: Non human primates 
免疫原	Recombinant full length protein corresponding to Human CREB aa 1-341. Sequence:

```
MTMESGAENQ QSGDAAVTEA ENQQMTVQAQ
PQIATLAQVS MPAAHATSSA PTVTLVQLPN
GQTVQVHGVI QAAQPSVIQS PQVQTVQSSC
KDLKRLFSGT QISTIAESED SQESVDSVTD
SQKRREILSR RPSYRKILND LSSDAPGVPR
IEEEKSEET SAPAITTVTV PTPIYQTSSG
QYIAITQGGA IQLANNGTDG VQGLQTLTMT
NAAATQPGTT ILQYAQTTDG QQILVPSNQV
VVQAASGDVQ TYQIRTAPTS TIAPGVVMAS
SPALPTQPAE EAARKREVRL MKNREAAREC
RRKKKEYVKC LENRVAVLEN QNKTLIEELK
ALKDLYCHKS D
```

Database link: [P16220](#)

 [Run BLAST with](#)

 [Run BLAST with](#)

阳性对照

HeLa cells; Human bladder and rectal cancer tissues; Human tonsil tissue; Human Jurkat, NIH/3T3, C2C12, MDA-MB-231, T98G, HeLa, 293T, HepG2, A431, A549, K562, MCF7 and U2OS whole cell lysates; Mouse MEF, 3T3L1 and C2C12 whole cell lysates; Rat NRK whole cell lysate; Monkey COS7 whole cell lysate.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 99% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	LB9
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab178322于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/500.
WB	★★★★★ (1)	1/500. Detects a band of approximately 43 kDa (predicted molecular weight: 36 kDa).
IP		Use at an assay dependent concentration. Suggested use 2 µg.
IHC-P		1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标

功能	This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Implicated in synchronization of circadian rhythmicity.
疾病相关	Defects in CREB1 may be a cause of angiomatoid fibrous histiocytoma (AFH) [MIM:612160]. A distinct variant of malignant fibrous histiocytoma that typically occurs in children and adolescents and is manifest by nodular subcutaneous growth. Characteristic microscopic features include lobulated sheets of histiocyte-like cells intimately associated with areas of hemorrhage and cystic pseudovascular spaces, as well as a striking cuffing of inflammatory cells, mimicking a lymph node metastasis. Note=A chromosomal aberration involving CREB1 is found in a patient with angiomatoid fibrous histiocytoma. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type.

序列相似性

Belongs to the bZIP family.
Contains 1 bZIP domain.
Contains 1 KID (kinase-inducible) domain.

翻译后修饰


Stimulated by phosphorylation. Phosphorylation of both Ser-133 and Ser-142 in the SCN regulates the activity of CREB and participates in circadian rhythm generation. Phosphorylation of Ser-133 allows CREBBP binding (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR.

Sumoylated by SUMO1. Sumoylation on Lys-304, but not on Lys-285, is required for nuclear localization of this protein. Sumoylation is enhanced under hypoxia, promoting nuclear localization and stabilization.

细胞定位

Nucleus.

图片



Lane 1: Wild-type HAP1 cell lysate (20 μ g)


Lane 2: CREB knockout HAP1 cell lysate (20 μ g)

Lane 3: HeLa cell lysate (20 μ g)

Lane 4: A431 cell lysate (20 μ g)

Lanes 1 - 4: Merged signal (red and green). Green - ab178322 observed at 44 kDa. Red - loading control, **ab18058**, observed at 124 kDa.

ab178322 was shown to recognize CREB when CREB knockout samples were used, along with additional cross-reactive bands. Wild-type and CREB knockout samples were subjected to SDS-PAGE. Ab178322 and **ab18058** (loading control to Vinculin) were diluted at 1/250 and 1/10,000 dilution respectively and incubated overnight at 4C. Blots were developed with IRDye® 800CW Goat anti-Rabbit IgG (H + L) and IRDye® 680 Goat anti-Mouse IgG (H + L) secondary antibodies at 1/10,000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-CREB antibody [LB9] (ab178322)

All lanes : Anti-CREB antibody [LB9] (ab178322) at 1/500 dilution

Lane 1 : Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

Lane 2 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 3 : K-562 (Human chronic myelogenous leukemia lymphoblast cell line) whole cell lysate

Lane 4 : Hep G2 (Human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 5 : NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate

Lane 6 : C2C12 whole cell lysate

Lane 7 : U-2 OS (Human bone osteosarcoma epithelial cell line) whole cell lysate

Lane 8 : A549 (Human lung carcinoma cell line) whole cell lysate


Lane 9 : MDA-MB-231 (Human breast adenocarcinoma cell line) whole cell lysate

Lane 10 : T98G whole cell lysate

Lysates/proteins at 30 µg per lane.


Predicted band size: 36 kDa

Observed band size: 43 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CREB antibody [LB9] (ab178322)

Immunohistochemical analysis of deparaffinized Human tonsil tissue labeling CREB with ab178322 at 1/100 dilution followed by DAB staining.




Immunoprecipitation - Anti-CREB antibody [LB9]
(ab178322)

Immunoprecipitation analysis of CREB was performed on untreated 293T cells. The antigen:antibody complex was formed by binding 500µg whole cell lysate with 2µg ab178322 and captured on 50µl Protein A/G Plus Agarose.

Lane 1: 293T lysate

Lane 2: IP of 293T lysate

WB detection used ab178322 at 1/1000 followed with a goat anti-mouse-HRP secondary antibody at 1/20,000.



Western blot - Anti-CREB antibody [LB9]
(ab178322)

All lanes : Anti-CREB antibody [LB9] (ab178322) at 1/500 dilution

Lane 1 : HeLa whole cell lysate

Lane 2 : 293T whole cell lysate

Lane 3 : A431 whole cell lysate

Lane 4 : A549 whole cell lysate

Lane 5 : MEF whole cell lysate

Lane 6 : MCF7 whole cell lysate

Lane 7 : U2OS whole cell lysate

Lane 8 : K562 whole cell lysate

Lane 9 : COS7 whole cell lysate

Lane 10 : 3T3L1 whole cell lysate

Lane 11 : C2C12 whole cell lysate

Lane 12 : NRK whole cell lysate


Lysates/proteins at 25 µg per lane.

Secondary

All lanes : goat anti-mouse-HRP at 1/20000 dilution


Developed using the ECL technique.

Predicted band size: 36 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CREB antibody [LB9]
(ab178322)

Immunohistochemical analysis of deparaffinized Human bladder cancer (left) and rectal cancer (right) tissues labeling CREB with ab178322 at 1/200 dilution followed by DAB staining.



Immunofluorescent analysis of untreated HeLa cells (formalin-fixed, 0.1% Triton X-100 permeabilized) labeling CREB with ab178322 at 1/400 dilution (green) followed with DyLight 488 goat anti-mouse IgG secondary antibody at 1/400 dilution. Nuclei (blue) were stained with Hoechst 33342 dye.



Immunofluorescent analysis of untreated HeLa cells (formalin-fixed, 0.1% Triton X-100 permeabilized) labeling CREB with ab178322 at 1/500 dilution (green) followed with DyLight 488 goat anti-mouse IgG secondary antibody at 1/400 dilution. Nuclei (blue) were stained with Hoechst 33342 dye.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors