

Anti-Cyclin T2/CCNT2 antibody ab96133

2 References **4 图像**

概述

产品名称	Anti-Cyclin T2/CCNT2抗体
描述	兔多克隆抗体to Cyclin T2/CCNT2
宿主	Rabbit
经测试应用	适用于: WB, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Human
免疫原	Recombinant fragment corresponding to Human Cyclin T2/CCNT2 aa 497-707.
阳性对照	WB: 293T, A431, H1299, HeLa, HepG2, MOLT4 and Raji cell lysates ICC/IF: Hela cell IHC-P: H1299 Xenograft; mouse brain tissue.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	<p>pH: 7.00</p> <p>Preservative: 0.01% Thimerosal (merthiolate)</p> <p>Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)</p>
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab96133于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/500 - 1/3000. Predicted molecular weight: 81 kDa.
IHC-P		1/100 - 1/250.
ICC/IF		1/100 - 1/200.

靶标

功能	Regulatory subunit of the cyclin-dependent kinase pair (CDK9/cyclin T) complex, also called positive transcription elongation factor B (P-TEFB), which is proposed to facilitate the transition from abortive to production elongation by phosphorylating the CTD (carboxy-terminal domain) of the large subunit of RNA polymerase II (RNAP II).
组织特异性	Ubiquitously expressed.
序列相似性	Belongs to the cyclin family. Cyclin C subfamily.
细胞定位	Nucleus.


图片



ab96133, at a 1/100 dilution, staining Cyclin T2/CCNT2 in paraffin embedded mouse brain tissue by Immunohistochemistry.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cyclin T2/CCNT2 antibody (ab96133)




Western blot - Anti-Cyclin T2/CCNT2 antibody
(ab96133)

Anti-Cyclin T2/CCNT2 antibody (ab96133) at 1/1000 dilution +
HepG2 whole cell lysate at 30 µg


Predicted band size: 81 kDa

7.5% SDS PAGE



Immunocytochemistry/ Immunofluorescence - Anti-
Cyclin T2/CCNT2 antibody (ab96133)

ab96133, at a 1/200 dilution, staining Cyclin T2/CCNT2 in
paraformaldehyde fixed A431 cells by Immunofluorescence.
Lower image: merged with DNA probe.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cylin T2/CCNT2 antibody (ab96133)

ab96133, at a 1/100 dilution, staining Cylin T2/CCNT2 in paraffin embedded H1299 (human cell line) xenograft by Immunohistochemistry.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors