abcam

Product datasheet

Anti-Desmin antibody [DE-U-10] - Cytoskeleton Marker ab6322

★★★★ 6 Abreviews 52 References 3 图像

概述

产品名称 Anti-Desmin抗体[DE-U-10] - Cytoskeleton Marker

描述 小鼠单克隆抗体[DE-U-10] to Desmin - Cytoskeleton Marker

宿主 Mouse

经测试应用 适用于: IHC-P

不适用于: ICC/IF

种属反应性 与反应: Mouse. Rat. Human

免疫原 Full length native protein (purified) corresponding to Pig Desmin.

阳性对照 IHC-P: Human skeletal muscle, Mouse cardiac muscle and rat colon tissue.

常规说明

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or

conjugation for your experiments, please contact orders@abcam.com.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

件能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

存储溶液

Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine

纯度 Protein G purified

克隆 单克隆 克隆编号 **DE-U-10**

骨髓瘤 unknown

同种型 lgG1

轻链类型 unknown

应用

The Abpromise guarantee Abpromise™承诺保证使用ab6322于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	****(1)	Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

应用说明 Is unsuitable for ICC/IF.

靶标

功能 Desmin are class-Ill intermediate filaments found in muscle cells. In adult striated muscle they form

a fibrous network connecting myofibrils to each other and to the plasma membrane from the

periphery of the Z-line structures.

疾病相关 Defects in DES are the cause of myopathy myofibrillar desmin-related (MFM-DES)

[MIM:601419]; also known as desmin-related myopathy (DRM). A neuromuscular disorder characterized by skeletal muscle weakness associated with cardiac conduction blocks, arrhythmias, restrictive heart failure, and by myofibrillar destruction with intracytoplasmic

accumulation of desmin-reactive deposits in cardiac and skeletal muscle cells.

Defects in DES are the cause of cardiomyopathy dilated type 1I (CMD1I) [MIM:604765]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function,

resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.

Defects in DES are the cause of neurogenic scapuloperoneal syndrome Kaeser type (Kaeser syndrome) [MIM:181400]. Kaeser syndrome is an autosomal dominant disorder with a peculiar

scapuloperoneal distribution of weakness and atrophy. A large clinical variability is observed ranging from scapuloperoneal, limb grindle and distal phenotypes with variable cardiac or respiratory involvement. Facial weakness, dysphagia and gynaecomastia are frequent additional symptoms. Affected men seemingly bear a higher risk of sudden, cardiac death as compared to affected women. Histological and immunohistochemical examination of muscle biopsy specimens


reveal a wide spectrum of findings ranging from near normal or unspecific pathology to typical,

myofibrillar changes with accumulation of desmin.

序列相似性 Belongs to the intermediate filament family.

细**胞定位** Cytoplasm.


图片



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Desmin antibody [DE-U-10] - Cytoskeleton Marker (ab6322)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse cardiac muscle tissue labelling Desmin with ab6322 at a dulution of 1/200. Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0). A ready to use HRP-conjugated goat anti-rabbit lgG H&L was used as the secondary antibody. Counter stained with Hematoxylin.

Image shows cytomplasmic staining on mouse cardiac muscle.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Desmin antibody [DE-U-10] - Cytoskeleton Marker (ab6322)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat colon tissue labelling Desmin with ab6322 at a dulution of 1/200. Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). A ready to use HRP-conjugated goat anti-rabbit lgG H&L was used as the secondary antibody. Counter stained with Hematoxylin.

Image shows cytomplasmic staining on smooth muscle of rat colon.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Desmin antibody [DE-U-10] - Cytoskeleton Marker (ab6322)

IHC image of Desmin staining in human skeletal muscle formalin fixed paraffin embedded tissue section, performed on a Leica BondTM system using the standard protocol F. The section was pretreated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab6322, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors