

Anti-DNA Ligase IV/LIG4 antibody [EPR16531] ab193353

敲除验证
重组
RabMAb

★★★★☆
4 Abreviews
13 References
6 图像

概述

产品名称	Anti-DNA Ligase IV/LIG4抗体[EPR16531]
描述	兔单克隆抗体[EPR16531] to DNA Ligase IV/LIG4
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, WB
种属反应性	与反应: Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HeLa, Jurkat and Ramos cell lysate. IHC-P Human thymus tissue. ICC/IF: SH-SY5Y and wild-type HAP1 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS, 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR16531
同种型	IgG

应用


The Abpromise guarantee **Abpromise™**承诺保证使用ab193353于以下的经测试应用
“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		1/50. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		Use a concentration of 5 µg/ml.
WB	★★★★☆ (4)	1/1000 - 1/2000. Predicted molecular weight: 103 kDa.

靶标

功能	Efficiently joins single-strand breaks in a double-stranded polydeoxynucleotide in an ATP-dependent reaction. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The LIG4-XRCC4 complex is responsible for the NHEJ ligation step, and XRCC4 enhances the joining activity of LIG4. Binding of the LIG4-XRCC4 complex to DNA ends is dependent on the assembly of the DNA-dependent protein kinase complex DNA-PK to these DNA ends.
组织特异性	Testis, thymus, prostate and heart.
疾病相关	LIG4 syndrome Severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-negative/NK-cell-positive with sensitivity to ionizing radiation
序列相似性	Belongs to the ATP-dependent DNA ligase family. Contains 2 BRCT domains.
细胞定位	Nucleus.

图片



Western blot - Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: DNA Ligase IV/LIG4 knockout HAP1 cell lysate (20 µg)

Lane 3: HepG2 cell lysate (20 µg)

Lane 4: HeLa cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab193353 observed at 103 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab193353 was shown to specifically react with DNA Ligase IV/LIG4 when DNA ligase IV knockout samples were used. Wild-type and DNA Ligase IV/LIG4 knockout samples were subjected to SDS-PAGE. ab193353 and **ab8245** (loading control to GAPDH) were diluted 1/1000 and 1/2000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353)

ab193353 staining DNA Ligase IV/LIG4 in wild-type HAP1 cells (top panel) and LIG4 knockout HAP1 cells (bottom panel). The cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab193353 at 5µg/ml dilution and **ab195889** at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.



Western blot - Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353)

All lanes : Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353) at 1/1000 dilution

Lane 1 : HeLa cell lysate

Lane 2 : Jurkat cell lysate


Lane 3 : Ramos cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution


Predicted band size: 103 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353)

Immunohistochemical analysis of paraffin embedded Human thymus tissue sections labeling DNA Ligase IV/LIG4 using ab193353 at a 1/50 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunofluorescent analysis of 4% paraformaldehyde fixed SH-SY5Y cells labeling DNA Ligase IV/LIG4 using ab193353 at a 1/50 dilution. A Goat anti rabbit IgG (Alexa Fluor®488) ([ab150077](#)) was used as the secondary at a 1/400 dilution. Counterstain DAPI. Cells were permeabilized using 0.1% Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353)

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-DNA Ligase IV/LIG4 antibody [EPR16531] (ab193353)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors