


### Anti-Factor X antibody ab97632

#### 1 图像

#### 概述

产品名称	Anti-Factor X抗体
描述	兔多克隆抗体to Factor X
宿主	Rabbit
经测试应用	适用于: WB
种属反应性	与反应: Human 预测可用于: Monkey 
免疫原	Recombinant protein fragment corresponding to a region within amino acids 33 and 282 of Human Factor X (NP_000495).
阳性对照	HeLa and HepG2 cell lysates
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.00</p> <p>Preservative: 0.01% Thimerosal (merthiolate)</p> <p>Constituents: 1.21% Tris, 0.75% Glycine, 10% Glycerol (glycerin, glycerine)</p>
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

#### 应用

The Abpromise guarantee      Abpromise™承诺保证使用ab97632于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/500 - 1/3000. Predicted molecular weight: 55 kDa.

靶标

功能	Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.
组织特异性	Plasma; synthesized in the liver.
疾病相关	Defects in F10 are the cause of factor X deficiency (FA10D) [MIM:227600]. A hemorrhagic disease with variable presentation. Affected individuals can manifest prolonged nasal and mucosal hemorrhage, menorrhagia, hematuria, and occasionally hemarthrosis. Some patients do not have clinical bleeding diathesis.
序列相似性	Belongs to the peptidase S1 family. Contains 2 EGF-like domains. Contains 1 Gla (gamma-carboxy-glutamate) domain. Contains 1 peptidase S1 domain.
翻译后修饰	The vitamin K-dependent, enzymatic carboxylation of some glutamate residues allows the modified protein to bind calcium. N- and O-glycosylated. The activation peptide is cleaved by factor IXa (in the intrinsic pathway), or by factor VIIa (in the extrinsic pathway). The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains.
细胞定位	Secreted.
形式	Cleaved into the following 3 chains: 1. Factor X light chain 2. Factor X heavy chain 3. Activated factor Xa heavy chain

图片



Anti-Factor X antibody (ab97632) at 1/5000 dilution + HepG2 whole cell lysate at 30 µg

**Predicted band size:** 55 kDa

10% SDS PAGE

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors