abcam

Product datasheet

Anti-FGFR2 antibody [EPR5180] ab109372





重组 RabMAb

★★★★★ 1 Abreviews 11 References 4 图像

概述

产品名称 Anti-FGFR2抗体[EPR5180]

描述 兔单克隆抗体[EPR5180] to FGFR2

宿主 Rabbit

经测试应用 适用于: WB, IP

不适用于: Flow Cyt,ICC/IF or IHC-P

种属反应性 与反应: Human

预测可用于: Mouse, Rat 📤

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 WB: MCF7, Jurkat, HeLa, K562, and T47-D cell lysates; IP: T-47D whole cell lysate.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.

存储溶液 pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.5% BSA

纯度 Protein A purified

克隆 单克隆 克隆编号 **EPR5180**

同种型 lgG

The Abpromise guarantee

Abpromise™承诺保证使用ab109372于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★☆ (1)	1/1000 - 1/10000. Detects a band of approximately 145 kDa (predicted molecular weight: 92 kDa).
IP		1/10 - 1/100.

应用说明

Is unsuitable for Flow Cyt,ICC/IF or IHC-P.

靶标

功能

疾病相关

Receptor for acidic and basic fibroblast growth factors.

Defects in FGFR2 are the cause of Crouzon syndrome (CS) [MIM:123500]; also called craniofacial dysostosis type I (CFD1). CS is an autosomal dominant syndrome characterized by craniosynostosis (premature fusion of the skull sutures), hypertelorism, exophthalmos and external strabismus, parrot-beaked nose, short upper lip, hypoplastic maxilla, and a relative mandibular prognathism.

Defects in FGFR2 are a cause of Jackson-Weiss syndrome (JWS) [MIM:123150]. JWS is an autosomal dominant craniosynostosis syndrome characterized by craniofacial abnormalities and abnormality of the feet: broad great toes with medial deviation and tarsal-metatarsal coalescence. Defects in FGFR2 are a cause of Apert syndrome (APRS) [MIM:101200]; also known as acrocephalosyndactyly type 1 (ACS1). APRS is a syndrome characterized by facio-cranio-synostosis, osseous and membranous syndactyly of the four extremities, and midface hypoplasia. The craniosynostosis is bicoronal and results in acrocephaly of brachysphenocephalic type. Syndactyly of the fingers and toes may be total (mitten hands and sock feet) or partial affecting the second, third, and fourth digits. Intellectual deficit is frequent and often severe, usually being associated with cerebral malformations.

Defects in FGFR2 are a cause of Pfeiffer syndrome (PS) [MIM:101600]; also known as acrocephalosyndactyly type V (ACS5). PS is characterized by craniosynostosis (premature fusion of the skull sutures) with deviation and enlargement of the thumbs and great toes, brachymesophalangy, with phalangeal ankylosis and a varying degree of soft tissue syndactyly. Three subtypes of Pfeiffer syndrome have been described: mild autosomal dominant form (type 1); cloverleaf skull, elbow ankylosis, early death, sporadic (type 2); craniosynostosis, early demise, sporadic (type 3).

Defects in FGFR2 are the cause of Beare-Stevenson cutis gyrata syndrome (BSCGS) [MIM:123790]. BSCGS is an autosomal dominant condition is characterized by the furrowed skin disorder of cutis gyrata, acanthosis nigricans, craniosynostosis, craniofacial dysmorphism, digital anomalies, umbilical and anogenital abnormalities and early death.

Defects in FGFR2 are the cause of familial scaphocephaly syndrome (FSPC) [MIM:609579]; also known as scaphocephaly with maxillary retrusion and mental retardation. FSPC is an autosomal dominant craniosynostosis syndrome characterized by scaphocephaly, macrocephaly, hypertelorism, maxillary retrusion, and mild intellectual disability. Scaphocephaly is the most common of the craniosynostosis conditions and is characterized by a long, narrow head. It is due to premature fusion of the sagittal suture or from external deformation.

Defects in FGFR2 are a cause of lacrimo-auriculo-dento-digital syndrome (LADDS) [MIM:149730]; also known as Levy-Hollister syndrome. LADDS is a form of ectodermal dysplasia, a heterogeneous group of disorders due to abnormal development of two or more ectodermal structures. LADDS is an autosomal dominant syndrome characterized by aplastic/hypoplastic lacrimal and salivary glands and ducts, cup-shaped ears, hearing loss, hypodontia and enamel hypoplasia, and distal limb segments anomalies. In addition to these cardinal features, facial dysmorphism, malformations of the kidney and respiratory system and abnormal genitalia have been reported. Craniosynostosis and severe syndactyly are not observed.

Defects in FGFR2 are the cause of Antley-Bixler syndrome (ABS) [MIM:207410]. ABS is a multiple congenital anomaly syndrome characterized by craniosynostosis, radiohumeral synostosis, midface hypoplasia, malformed ears, arachnodactyly and multiple joint contractures.

ABS is a heterogeneous disorder and occurs with and without abnormal genitalia in both sexes.

序列相似性

Belongs to the protein kinase superfamily. Tyr protein kinase family. Fibroblast growth factor receptor subfamily.


Contains 3 lg-like C2-type (immunoglobulin-like) domains.

Contains 1 protein kinase domain.

细胞定位

Secreted and Cell membrane.

图片



Western blot - Anti-FGFR2 antibody [EPR5180] (ab109372)

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: FGFR2 knockout HAP1 cell lysate (20 µg)


Lane 3: HeLa cell lysate (20 µg)

Lane 4: Jurkat cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab109372 observed at 160 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab109372 was shown to specifically react with FGFR2 when FGFR2 knockout samples were used. Wild-type and FGFR2 knockout samples were subjected to SDS-PAGE.

Ab109372 and ab8245 (loading control to GAPDH) were diluted at 1/1000 and 1/10,000 dilution respectively and incubated overnight at 4C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10,000 dilution for 1 hour at room temperature before imaging.




(ab109372)

All lanes : Anti-FGFR2 antibody [EPR5180] (ab109372) at 1/1000 dilution

Lane 1 : MCF7 cell lysate
Lane 2 : Jurkat cell lysate
Lane 3 : HeLa cell lysate
Lane 4 : K562 cell lysate
Lane 5 : T47-D cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 92 kDa **Observed band size:** 145 kDa



Immunoprecipitation - Anti-FGFR2 antibody [EPR5180] (ab109372)

Purified ab109372 at 1/40 dilution ($2\mu g$) immunoprecipitating FGFR2 in T-47D whole cell lysate.

Lane 1 (input): T-47D (Human ductal breast epithelial tumor epithelial cell) whole cell lysate 10µg


Lane 2 (+): ab109372 + T-47D whole cell lysate.

Lane 3 (-): Rabbit monoclonal $\lg G$ (ab172730) instead of ab109372 in T-47D whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) (<u>ab131366</u>) (1/1000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST. Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: 145 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors