

### FITC Anti-GFP antibody ab6662

★★★★★ [15 Abreviews](#) [171 References](#) [5 图像](#)

#### 概述

产品名称	FITC荧光Anti-GFP抗体
描述	FITC荧光山羊多克隆抗体to GFP
宿主	Goat
偶联物	FITC. Ex: 493nm, Em: 528nm
经测试应用	<b>适用于:</b> IHC-FoFr, IHC-Fr, WB, ICC/IF
种属反应性	<b>与反应:</b> Species independent
免疫原	Recombinant full length protein corresponding to GFP aa 1-246. Sequence:  MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKL TLKFICTT GKLPPWPPTLVTTFSYGVQCFSRYPDHMKQHDFKSAPEGY VQERTIFF KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLE YNYNSHNV YIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQNTPIGDGP VLLPDNHY LSTQSALSKDPNEKRDHMLLEFVTAAGITHGMDELYK

Database link: [P42212](#)

 [Run BLAST with](#)

 [Run BLAST with](#)

**阳性对照** WB: Recombinant A. victoria GFP protein.

**常规说明** Designed to detect GFP and its variants in ELISA (sandwich or capture), immunoblotting and immunoprecipitation. Fluorescein conjugated anti-GFP was assayed by immunofluorescence microscopy on prokaryotic (*E.coli*) and eukaryotic (CHO cells) expression systems and was shown to detect GFP containing inserts. Significant amplification of signal was detected using fluorochrome conjugated anti-GFP relative to the fluorescence of GFP alone.  
In case of unexpected background, use pre-adsorbed secondary antibodies.

Fluorescein isothiocyanate (FITC) (MW 390 daltons) Absorption Wavelength: 495 nm Emission Wavelength: 528 nm Fluorochrome/Protein Ratio: 3.5 moles FITC per mole of Goat IgG

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 6.50 Preservative: 0.01% Sodium azide Constituents: 0.42% Potassium phosphate, 0.87% Sodium chloride, 1% BSA  BSA Immunoglobulin and Protease free
纯度	Protein G purified
纯化说明	GFP Fluorescein Conjugated Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Green Fluorescent Protein ( <i>Aequorea victoria</i> ) coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
Primary antibody说明	Designed to detect GFP and its variants in ELISA (sandwich or capture), immunoblotting and immunoprecipitation. Fluorescein conjugated anti-GFP was assayed by immunofluorescence microscopy on prokaryotic ( <i>E.coli</i> ) and eukaryotic (CHO cells) expression systems and was shown to detect GFP containing inserts. Significant amplification of signal was detected using fluorochrome conjugated anti-GFP relative to the fluorescence of GFP alone.
克隆	多克隆
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise™** 承诺保证使用ab6662于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-FoFr	★★★★★ (1)	1/250.
IHC-Fr	★★★★★ (5)	Use at an assay dependent concentration.
WB	★★★★☆ (1)	1/10000.
ICC/IF	★★★★☆ (4)	1/500 - 1/2500.

## 靶标

相关性	<b>Function:</b> Energy-transfer acceptor. Its role is to transduce the blue chemiluminescence of the protein aequorin into green fluorescent light by energy transfer. Fluoresces in vivo upon receiving energy from the Ca <sup>2+</sup> -activated photoprotein aequorin.
-----	--

**Subunit structure:** Monomer.

**Tissue specificity:** Photocytes.

**Post-translational modification:** Contains a chromophore consisting of modified amino acid residues. The chromophore is formed by autocatalytic backbone condensation between Ser-65 and Gly-67, and oxidation of Tyr-66 to didehydrotyrosine. Maturation of the chromophore requires nothing other than molecular oxygen.


**Biotechnological use:** Green fluorescent protein has been engineered to produce a vast number of variously colored mutants, fusion proteins, and biosensors. Fluorescent proteins and its mutated allelic forms, blue, cyan and yellow have become a useful and ubiquitous tool for making chimeric proteins, where they function as a fluorescent protein tag. Typically they tolerate N- and C-terminal fusion to a broad variety of proteins. They have been expressed in most known cell types and are used as a noninvasive fluorescent marker in living cells and organisms. They enable a wide range of applications where they have functioned as a cell lineage tracer, reporter of gene expression, or as a measure of protein-protein interactions. Can also be used as a molecular thermometer, allowing accurate temperature measurements in fluids. The measurement process relies on the detection of the blinking of GFP using fluorescence correlation spectroscopy.

**Sequence similarities:** Belongs to the GFP family.

**Biophysicochemical properties:** Absorption: Abs(max)=395 nm

Exhibits a smaller absorbance peak at 470 nm. The fluorescence emission spectrum peaks at 509 nm with a shoulder at 540 nm.

## 图片




FITC Anti-GFP antibody (ab6662) + Recombinant A. victoria GFP protein ([ab84191](#))

### Secondary

Fluorescein goat secondary antibody for 60 minutes at RT at 1/1000 dilution

**Block:** Blocking buffer for 30 minutes at RT.



Immunohistochemistry (Frozen sections) - FITC


Anti-GFP antibody (ab6662)

Kim et al PLoS Genet. 2018 Feb 8;14(2):e1007204. doi: 10.1371/journal.pgen.1007204. eCollection 2018 Feb. Fig 4. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

### Oscillation of Cyclin E and E2F target gene expression is deregulated in *de2f1b* mutant salivary glands.

Salivary glands of control and *de2f1b* mutant early (80–85 hr AEL) third instar larvae expressing PCNA-GFP (green, ab6662) are stained with anti-dE2F1 (red). The region where high PCNA-GFP is observed with low dE2F1 is marked by an asterisk.

For Immunostaining, third instar imaginal discs and salivary glands were dissected in PBS and immediately fixed in 4% formaldehyde in PBS for 20 minutes at room temperature with the exception of tissues subjected to anti-dE2F1 staining that were fixed for 30 minutes on ice. Fixed tissues were then washed with 0.3% PBST (0.3% TritonX-100 in 1XPBS) and 0.1% PBST (0.1% TritonX-100 in 1XPBS). Samples were incubated with appropriate amount of primary antibody in 0.1% PBST and 1%BSA overnight. Samples were then washed with 0.1% PBST, incubated in secondary antibody in 0.1% PBST and 1% BSA for 2 hours, followed by several washes in 0.1% PBST prior to mounting.



Immunohistochemistry (Frozen sections) - FITC

Anti-GFP antibody (ab6662)


This image is courtesy of an Abreview submitted by Dr Joshua Breunig

ab6662 staining mouse brain tissue sections (inducible GFP reporter) by IHC-Fr.

The tissue was paraformaldehyde fixed and blocked with serum and then incubated with the antibody at a 1/1000 dilution for 1 hour.

Staining is shown in the left hand panel. The middle panel shows staining with a rabbit anti-GFP antibody and the right hand panel shows the merged images (plus DAPI).

ab6662 gives no noticable background and it is found that when viewing on an epifluorescent the exposure time is significantly reduced.



Immunohistochemistry (Frozen sections) - FITC  
Anti-GFP antibody (ab6662)

### Immunofluorescence Microscopy using ab6662.

**Tissue:** Sf-1:Cre mice crossed to the Z/EG reporter line. Mouse brain (coronal view, 20X magnification).

**Fixation:** 4% PFA/PBS with o/n fixation, and subsequently transferred to a 30% sucrose solution.


**Antigen retrieval:** Frozen in OCT freezing medium (Sakura) and cryostat sectioned at 40 microns.

Goat anti-GFP was used at 1/500 dilution in free floating immunohistochemistry to detect GFP.

**Secondary antibody:** Fluorochrome conjugated Anti-goat IgG secondary antibody was used for detection at 1:500 at 1/10,000 for 45 minutes at RT.

**Localization:** Sf-1+ neurons and their processes of the ventromedial nucleus of the hypothalamus.

**Staining:** eGFP as green fluorescent signal and sections were counterstained with DAPI.



Immunohistochemistry (Frozen sections) - FITC  
Anti-GFP antibody (ab6662)

These pictures show confocal immunofluorescence using GFP-expressing glial cells (green) transplanted into the lesioned rat spinal cord.

Detected using ab6662 and a standard FITC filter set. Axons are labeled red by an antibody to neurofilament-200 and a rhodamine secondary antibody. The upper panel shows the centre of the transplant site at low power. Numerous GFP-positive cells can be seen mingling with axons. The lower panel shows, at high power in a single optical section, how ab6662 reveals the morphology of the transplanted cells to such an extent that their close interactions with axons are obvious - the cell depicted can be seen wrapping around a neurofilament-200 positive axon.

These images were kindly supplied as part of the review submitted by Andrew Toft.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors