abcam

Product datasheet

Anti-Galectin 1 antibody [EPR3205] - Low endotoxin, Azide free ab225540



RabMAb

4 References 2 图像

概述

常规说明

产品名称 Anti-Galectin 1抗体[EPR3205] - Low endotoxin, Azide free

描述 兔单克隆抗体[EPR3205] to Galectin 1 - Low endotoxin, Azide free

宿主 Rabbit

经测试应用 适用于: WB, IHC-P, ICC/IF

种属反应性 与反应: Mouse, Rat, Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 WB: HL60, K562, and NIH/3T3 cell lysates. Mouse heart, kidney, and spleen tissue lysates. Rat heart and kidney tissue lysates; ICC/IF: NIH/3T3 and K562 cells. IHC-P: Human placenta tissue.

ab225540 is the carrier-free version of ab108389.

Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our <u>conjugation kits</u> for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**[®] **patents**.

1

Our <u>Low endotoxin, azide-free formats</u> have low endotoxin level (≤ 1 EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C. Do Not Freeze.

存储溶液 pH: 7.20

Constituent: PBS

无载体 是

纯**度** Protein A purified

同种型 lgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab225540于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.

벎	Ц	1	k.	ê
48	-	4	7	Λ

功能 May regulate apoptosis, cell proliferation and cell differentiation. Binds beta-galactoside and a

wide array of complex carbohydrates. Inhibits CD45 protein phosphatase activity and therefore

the dephosphorylation of Lyn kinase.


组织**特异性** Expressed in placenta, maternal decidua and fetal membranes. Within placenta, expressed in

trophoblasts, stromal cells, villous endothelium, syncytiotrophoblast apical membrane and villous stroma. Within fetal membranes, expressed in amnion, chorioamniotic mesenchyma and chorion (at protein level). Expressed in cardiac, smooth, and skeletal muscle, neurons, thymus, kidney and

hematopoietic cells.

序列相似性 Contains 1 galectin domain.

细胞定位 Secreted > extracellular space > extracellular matrix.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Galectin 1 antibody

[EPR3205] - Low endotoxin, Azide free (ab225540)

Immunohistochemical analysis of paraffin-embedded human placenta tissue labelling Galectin 1 with $\underline{ab108389}$ at 1/3000 dilution (0.42 µg/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB ($\underline{ab209101}$) secondary antibody. Positive staining on vessels in human placenta tissue. The section was incubated with $\underline{ab108389}$ for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution1) for 20 mins.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab108389</u>).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors