

# Anti-gamma H2A.X (phospho S139) antibody [3F2] ab22551

★★★★★ [13 Abreviews](#) [150 References](#) [8 图像](#)

## 概述

产品名称	Anti-gamma H2A.X (phospho S139)抗体[3F2]
描述	小鼠单克隆抗体[3F2] to gamma H2A.X (phospho S139)
宿主	Mouse
经测试应用	适用于: WB, IHC-P, ICC/IF, Flow Cyt
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide corresponding to Human gamma H2A.X (phospho S139). Synthetic peptide sequence surrounding phosphorylated Ser139
阳性对照	WB: Jurkat (treated with staurosporin) cell lysate. ICC: HepG2, A549 and HeLa cells IHC-P: Human breast tissue; Postnatal mouse lung sections with DNA damage in airway cells. Flow Cyt: HeLa cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.05% Sodium azide Constituents: PBS, 0.1% BSA
纯度	Protein G purified
克隆	单克隆
克隆编号	3F2
同种型	IgG1
轻链类型	kappa

应用

**The Abpromise guarantee**      **Abpromise™**承诺保证使用ab22551于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★☆ (4)	Use a concentration of 1 µg/ml. Detects a band of approximately 17 kDa.
IHC-P	★★★★☆ (2)	Use at an assay dependent concentration.
ICC/IF	★★★★☆ (7)	Use a concentration of 2 - 4 µg/ml.
Flow Cyt		Use 1µg for 10 <sup>6</sup> cells. <b>ab170190</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

靶标


功能	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.
序列相似性	Belongs to the histone H2A family.
发展阶段	Synthesized in G1 as well as in S-phase.
结构域	The [ST]-Q motif constitutes a recognition sequence for kinases from the PI3/PI4-kinase family.
翻译后修饰	Phosphorylated on Ser-140 (to form gamma-H2AFX or H2AX139ph) in response to DNA double strand breaks (DSBs) generated by exogenous genotoxic agents and by stalled replication forks, and may also occur during meiotic recombination events and immunoglobulin class switching in lymphocytes. Phosphorylation can extend up to several thousand nucleosomes from the actual site of the DSB and may mark the surrounding chromatin for recruitment of proteins required for DNA damage signaling and repair. Widespread phosphorylation may also serve to amplify the damage signal or aid repair of persistent lesions. Phosphorylation of Ser-140 (H2AX139ph) in response to ionizing radiation is mediated by both ATM and PRKDC while defects in DNA replication induce Ser-140 phosphorylation (H2AX139ph) subsequent to activation of ATR and PRKDC. Dephosphorylation of Ser-140 by PP2A is required for DNA DSB repair. In meiosis, Ser-140 phosphorylation (H2AX139ph) may occur at synaptonemal complexes during leptotene as an ATM-dependent response to the formation of programmed DSBs by SPO11. Ser-140 phosphorylation (H2AX139ph) may subsequently occurs at unsynapsed regions of both autosomes and the XY bivalent during zygotene, downstream of ATR and BRCA1 activation. Ser-140 phosphorylation (H2AX139ph) may also be required for transcriptional repression of unsynapsed chromatin and meiotic sex chromosome inactivation (MSCI), whereby the X and Y chromosomes condense in pachytene to form the heterochromatic XY-body. During immunoglobulin class switch recombination in lymphocytes, Ser-140 phosphorylation

(H2AX139ph) may occur at sites of DNA-recombination subsequent to activation of the activation-induced cytidine deaminase AICDA. Phosphorylation at Tyr-143 (H2AXY142ph) by BAZ1B/WSTF determines the relative recruitment of either DNA repair or pro-apoptotic factors. Phosphorylation at Tyr-143 (H2AXY142ph) favors the recruitment of APBB1/FE65 and pro-apoptosis factors such as MAPK8/JNK1, triggering apoptosis. In contrast, dephosphorylation of Tyr-143 by EYA proteins (EYA1, EYA2, EYA3 or EYA4) favors the recruitment of MDC1-containing DNA repair complexes to the tail of phosphorylated Ser-140 (H2AX139ph). Monoubiquitination of Lys-120 (H2AXK119ub) by RING1 and RNF2/RING2 complex gives a specific tag for epigenetic transcriptional repression. Following DNA double-strand breaks (DSBs), it is ubiquitinated through 'Lys-63' linkage of ubiquitin moieties by the E2 ligase UBE2N and the E3 ligases RNF8 and RNF168, leading to the recruitment of repair proteins to sites of DNA damage. Monoubiquitination and ionizing radiation-induced 'Lys-63'-linked ubiquitination are distinct events.

## 细胞定位


Nucleus. Chromosome.

## 图片




ICC/IF image of ab22551 stained HepG2 cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab22551, 10µg/ml) overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti-mouse IgG - H&L, pre-adsorbed (**ab96879**) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Immunocytochemistry/ Immunofluorescence - Anti-gamma H2A.X(phospho S139) antibody [3F2] (ab22551)



Western blot - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)


Western blot analysis of Phospho-H2A.X (phospho S139) (ab22551) at a concentration of 1 µg/mL on Jurkat cell untreated (Lane 1) and Jurkat cell stimulated with staurosporine (Lane 2) followed by HRP conjugated goat anti-mouse IgG (H+L) Secondary Antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)


IHC image of ab22551 staining in Human breast formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab22551, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Flow Cytometry - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)


Overlay histogram showing HeLa cells stained with ab22551 (red line). The cells were fixed with 100% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab22551, 1µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was Mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)


Immunohistochemistry analysis of phospho-Histone H2A.X pSer140 labelled with ab22551 in postnatal mouse (PN19) lung sections. Antigen retrieval from 4% PFA, paraffin embedded sections was performed using heat induced epitope retrieval (HIER) method with sodium citrate buffer (pH 6.0). Following antigen retrieval, tissues were blocked and probed with ab22551 at a dilution of 1:400. Increased staining intensity was observed in a genetic mouse model with DNA damage in airway cells.

Left: Control; Right: A genetic model with DNA damage in airway cells




Immunocytochemistry/ Immunofluorescence - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)

Immunofluorescence staining of Phospho-H2AX with ab22551 at 4 ug/ml in A549 cells. Cells were treated with vehicle (control; 0.1% DMSO in media) (right) or with 50 µM etoposide for 1 hour (left).



Immunocytochemistry/ Immunofluorescence - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)

ab22551 labelling gamma H2A.X (phospho S139) in HeLa cells by immunocytochemistry/immunofluorescence.



A431 cell nucleus (DAPI signal) showing three gamma-H2AX foci structures.

Primary antibody H2A.X (phospho S139) antibody [3F2], ab22551, 100ug.

Secondary antibody Mouse IgG-Fc (FITC), **ab97264**, 1mg.

Immunocytochemistry/ Immunofluorescence - Anti-gamma H2A.X (phospho S139) antibody [3F2] (ab22551)

This image is courtesy of Jorge E. Gonzalez (CPHR, La Habana, Cuba), Joan F. Barquinero (UAB, Barcelona, Spain) and Jessica Martínez (UAB, Universitat Autònoma de Barcelona, Spain).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors