

Anti-GAPDH antibody [EPR16891] - BSA and Azide free ab199553

重组 RabMAb

★★★★★ [1 Abreviews](#) [3 References](#) [12 图像](#)

概述

产品名称	Anti-GAPDH抗体[EPR16891] - BSA and Azide free
描述	兔单克隆抗体[EPR16891] to GAPDH - BSA and Azide free
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), IP, IHC-P, WB, ICC/IF
种属反应性	与反应: Mouse, Rat, Chicken, Human, Zebrafish, African green monkey, Xenopus tropicalis 预测可用于: Rabbit, Fish 
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HeLa, UMNSAH/DF-1, Jurkat, COS-1, RAW 264.7 and PC-12 whole cell lysates; Human fetal brain and heart lysates; Xenopus(X. tropicalis) muscle lysate; Zebrafish lysate; Mouse kidney and spleen lysates; Rat brain lysate. IHC-P: Human transitional cell carcinoma of bladder, Mouse spleen and Rat spleen tissues. ICC/IF: HeLa cells. Flow: Jurkat cells. IP: HeLa whole cell extract
常规说明	<p>ab199553 is the carrier-free version of ab181602.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR16891
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab199553 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB	★★★★★ (1)	Use at an assay dependent concentration. Detects a band of approximately 36 kDa (predicted molecular weight: 36 kDa).
ICC/IF		Use at an assay dependent concentration.

靶标

功能 Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-

phosphate (G3P) into 3-phospho-D-glyceroyl phosphate.

Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1/5.

Belongs to the glyceraldehyde-3-phosphate dehydrogenase family.

S-nitrosylation of Cys-152 leads to interaction with SIAH1, followed by translocation to the nucleus.

ISGylated.

Cytoplasm > cytosol. Nucleus. Cytoplasm > perinuclear region. Membrane. Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions.


通路

序列相似性

翻译后修饰

细胞定位

图片



Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553) at 1/100000 dilution + HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysates at 15 µg


Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 36 kDa


Blocking and Diluting buffer and concentration: 5% NFDm/TBST

Exposure time: 3 seconds



Immunocytochemistry/immunofluorescence staining of 4% paraformaldehyde fixed; 0.1% triton X 100 permeabilized HeLa (human cervix adenocarcinoma) cells labeling GAPDH with [ab181602](#) at dilution of 1/500. The secondary antibody used was Alexa Fluor[®] 488; goat anti-rabbit IgG ([ab150077](#)) at a dilution of 1/400. Nucleus was counter-stained with DAPI (blue). [ab7291](#), a mouse anti-tubulin antibody (1/500) was used to stain tubulin along with [ab150120](#) (AlexaFluor[®]594 goat anti-mouse secondary, 1/500). The negative controls are shown in the bottom middle and right hand panels- for negative control 1 primary antibody ([ab181602](#); 1/500) and secondary antibody ([ab150120](#); 1/500) was used. For negative control 2 primary antibody ([ab7291](#); 1/500) and secondary antibody ([ab150077](#); 1/400) was used.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab181602](#)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Immunohistochemical analysis of paraffin-embedded human transitional cell carcinoma of bladder tissue labeling GAPDH with **ab181602** at 1/2000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasmic and nucleus staining on the tumor cells of transitional cell carcinoma of human bladder is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab181602**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling GAPDH with **ab181602** at 1/2000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Nucleus and cytoplasmic staining on lymphocytes of mouse spleen is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab181602**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Immunohistochemical analysis of paraffin-embedded rat spleen tissue labeling GAPDH with **ab181602** at 1/2000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Nucleus and cytoplasmic staining on lymphocyte of rat spleen is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab181602**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

This data was developed using **ab181602**, the same antibody clone in a different buffer formulation. Different batches of **ab181602** were tested on HeLa (Human cervix adenocarcinoma epithelial cell) lysate at 1.0 µg/ml. 15 µg of lysate was loaded in each lane. Bands observed at 36 kDa.




Immunocytochemistry/ Immunofluorescence - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Clone EPR16891 (ab199553) has been successfully conjugated by Abcam. This image was generated using Anti-GAPDH antibody [EPR16891] (Alexa Fluor® 488). Please refer to [ab201768](#) for protocol details.

[ab201768](#) staining GAPDH in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with [ab201768](#) at 1/200 dilution (shown in green) and [ab195889](#), Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at 2µg/ml (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Clone EPR16891 (ab199553) has been successfully conjugated by Abcam. This image was generated using Anti-GAPDH antibody [EPR16891] (Alexa Fluor® 647). Please refer to [ab201272](#) for protocol details.

[ab201272](#) staining GAPDH in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with [ab201272](#) at 1/100 dilution (shown in red) and [ab195887](#), Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 2µg/ml (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Flow Cytometry (Intracellular) - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)


Clone EPR16891 (ab199553) has been successfully conjugated by Abcam. This image was generated using Anti-GAPDH antibody [EPR16891] (PE). Please refer to [ab224004](#) for protocol details.

Overlay histogram showing HeLa cells stained with [ab224004](#) (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody ([ab224004](#), 1/1000 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was Rabbit IgG (monoclonal) Phycoerythrin ([ab209478](#)) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50 mW Yellow/Green laser (561nm) and 586/15 bandpass filter.


This antibody gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min)/permeabilized with 0.1% PBS-Triton X-100 for 15 min used under the same conditions.



Flow Cytometry (Intracellular) - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed Jurkat (Human T cell leukemia cells from peripheral blood) cells labeling GAPDH with [ab181602](#) at 1/180 dilution (red) compared with a rabbit monoclonal IgG isotype control (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab181602](#)).



Immunoprecipitation - Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

GAPDH was immunoprecipitated from 1mg of HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell extract with **ab181602** at 1/60 dilution. Western blot was performed from the immunoprecipitate using **ab181602** at 1/5000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution. **Lane 1:** HeLa whole cell extract.

Lane 2: PBS instead of HeLa whole cell extract.

Blocking and dilution buffer and concentration: 5% NFDN/TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab181602**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-GAPDH antibody [EPR16891] - BSA and Azide free (ab199553)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors