# abcam

## Product datasheet

## Anti-Geminin antibody [1A8] ab104306

★★★★★ 1 Abreviews 7 References 7 图像

#### 概述

产品名称 Anti-Geminin抗体[1A8]

描述 小鼠单克隆抗体[1A8] to Geminin

**宿主** Mouse

经测试应用 适用于: IHC-P, ICC/IF, WB, Flow Cyt

种属反应性 与反应: Human

免疫原 Recombinant fragment, corresponding to amino acids 110-209 of Human Geminin (AAH05185)

with a proprietary tag.

**阳性**对照 Partial tagged recombinant Geminin; Geminin transfected 293T cell lysate; Human lymph node;

Human placenta; HeLa cells.

常规说明 This product was changed from ascites to tissue culture supernatant on 22 May 2019. Please

note that the dilutions may need to be adjusted accordingly. If you have any questions, please do

not hesitate to contact our scientific support team.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

**存储溶液** pH: 7.40

Constituent: 100% PBS

纯**度** Tissue culture supernatant

纯**化**说明 Purified from TCS.

**克隆** 单克隆

**克隆编号** 1A8

1

**同种型** IgG1

轻链类型 kappa

#### 应用

## The Abpromise guarantee Abpromise™承诺保证使用ab104306于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF	**** <u>(1)</u>	Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Predicted molecular weight: 24 kDa.
Flow Cyt		Use at an assay dependent concentration. <u>ab170190</u> - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.

## 靶标

功能 Inhibits DNA replication by preventing the incorporation of MCM complex into prereplication

complex (pre-RC). It is degraded during the mitotic phase of the cell cycle. Its destruction at the


metaphase-anaphase transition permits replication in the succeeding cell cycle.

序列相似性 Belongs to the geminin family.

发展阶段 Absent during G1 phase, accumulates during S, G2, and M phases, and disappears at the time of

the metaphase-anaphase transition.

## 图片




Western blot - Anti-Geminin antibody [1A8] (ab104306)

Anti-Geminin antibody [1A8] (ab104306) at 5  $\mu$ g/ml + Immunogen at 0.2  $\mu$ g


Predicted band size: 24 kDa

The tagged immunogen has a predicted MWt of 37 kDa.

This image was generated using the ascites version of the product.



Western blot - Anti-Geminin antibody [1A8] (ab104306)




Lane 1: Geminin transfected 293T cell lysate

Lane 2: Non-transfected 293T cell lysate

Lysates/proteins at 25 µg per lane.

Predicted band size: 24 kDa


This image was generated using the ascites version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Geminin antibody [1A8] (ab104306)

ab104306, at 3  $\mu$ g/ml, staining Geminin in formalin-fixed paraffinembedded Human lymph node by Immunohistochemistry.


This image was generated using the ascites version of the product.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Geminin antibody [1A8] (ab104306)

ab104306, at 1µg/ml, staining Geminin in formalin-fixed paraffinembedded Human placenta by Immunohistochemistry.


This image was generated using the ascites version of the product.



Immunocytochemistry/ Immunofluorescence - Anti-Geminin antibody [1A8] (ab104306)

ab104306, at 10  $\mu$ g/ml, staining Geminin in HeLa cells by Immunofluorescence.


This image was generated using the ascites version of the product.



Flow Cytometry - Anti-Geminin antibody [1A8] (ab104306)

Overlay histogram showing HeLa cells stained with ab104306 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab104306,  $1\mu g/1x10^6$  cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse lgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse lgG1 [ICIGG1] (ab91353,  $2\mu g/1x10^6$  cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

This image was generated using the ascites version of the product.



Immunocytochemistry/ Immunofluorescence - Anti-Geminin antibody [1A8] (ab104306)

This image is courtesy of an Abreview submitted by Melody Jimenez

Immunofluorescence analysis of Caki-2 (Clear cell renal carcinoma) cells, staining Geminin with ab104306.

Cells were fixed with formaldehyde, permeabilized with 0.25% Triton X-100 and blocked with 1% BSA for 30 minutes at  $25^{\circ}$ C. Primary antibody was concentrated using **ab102778** Antibody Concentration Kit and conjugated to Rhodamine using **ab102915** EasyLink Rhodamine Conjugation Kit. Samples were incubated with conjugated primary antibody (20  $\mu$ g/ml in diluent) for 2 hours at 25°C.

This image was generated using the ascites version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors