



# Anti-Glucocorticoid Receptor alpha antibody ab3580

★★★★★ **3 Abreviews** **28 References** **9 图像**

### 概述

产品名称	Anti-Glucocorticoid Receptor alpha抗体
描述	兔多克隆抗体to Glucocorticoid Receptor alpha
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF
种属反应性	与反应: Human
免疫原	Synthetic peptide corresponding to Human Glucocorticoid Receptor alpha aa 755-771. Sequence: CEIITNQIPKYSNGNIKK  Database link: <b>P04150</b> (Peptide available as <b>ab39764</b> )   <b>Run BLAST with</b>  <b>Run BLAST with</b>
阳性对照	IHC: human cervical carcinoma, heart tissue, tonsil tissue; ICC: U251, A2058, HeLa
常规说明	GR alpha proteins has many isoforms e.g. GR alpha-A, Alpha-2, GR-A alpha, Alpha-B. Please check Uniprot database or PMID 15866175 for more information.  The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.  If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituent: 99% PBS
纯度	Immunogen affinity purified

克隆

多克隆

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab3580于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (1)	1/20.
ICC/IF	★★★☆☆ (1)	1/100 - 1/200.

靶标


相关性

Glucocorticoids are a family of steroids necessary for the regulation of energy metabolism and the immune and inflammatory responses. These compounds exert their effect through their interaction with the Glucocorticoid Receptor (GR) and that complex's subsequent association with DNA. All normal mammalian tissues examined to date have been shown to contain GR. The human GR exists in two forms, alpha and beta, which are thought to be the result of alternative splicing of a single gene. Sequence analysis indicates that alpha and beta forms of human GR are 777 and 742 amino acids long, respectively. They are identical up to residue 727, after which they diverge. After ligand binding, the 94 kDa GR alpha isoform translocates from the cytoplasm to the nucleus where it regulates gene expression. In contrast, the 90 kDa GR beta isoform does not appear to bind either glucocorticoid agonists or antagonists, and has been localized predominantly in the nucleus independent of hormone treatment in some human cell lines. Studies suggest that human GR alpha has a greater affinity for GR response elements (GREs) than GR beta only when in the ligand bound state.

细胞定位


Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand; nuclear after ligand-binding.

图片




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor alpha antibody - ChIP Grade (ab3580)

Immunocytochemistry/Immunofluorescence analysis of U-251 MG (Human brain glioma cell line) cells labeling Glucocorticoid Receptor alpha (green) with ab3580 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucocorticoid Receptor alpha antibody (ab3580)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized human tonsil tissue tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:50 with a rabbit polyclonal antibody recognizing Glucocorticoid Receptor alpha ab3580 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor alpha antibody - ChIP Grade (ab3580)

Immunocytochemistry/Immunofluorescence analysis of HeLa (Human epithelial adenocarcinoma cell line) cells labeling Glucocorticoid Receptor alpha (green) with ab3580 at 1/100. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucocorticoid Receptor alpha antibody (ab3580)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized human cervical carcinoma tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:20 with a rabbit polyclonal antibody recognizing Glucocorticoid Receptor alpha ab3580 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor alpha antibody - ChIP Grade (ab3580)

Immunocytochemistry/Immunofluorescence analysis of A2058 (Human metastatic melanoma cell line) cells labeling Glucocorticoid Receptor alpha (green) with ab3580 at 1/200. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. A DyLight 488-conjugated secondary antibody was used. 60X magnification. Right - negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucocorticoid Receptor alpha antibody (ab3580)


Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized human heart tissue tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a rabbit polyclonal antibody recognizing Glucocorticoid Receptor alpha ab3580 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.



Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor alpha antibody - ChIP Grade (ab3580)

ab3580 staining glucocorticoid receptor in serum starved HeLa (Human epithelial adenocarcinoma cell line) cells treated with rosiglitazone (**ab120762**), by ICC/IF. Changes in localization of glucocorticoid receptor (translocation from cytoplasm to nucleus) correlates with increased concentration of rosiglitazone, as described in literature.


The cells were incubated at 37°C for 1h in media containing different concentrations of **ab120762** (rosiglitazone) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab3580 (5 μg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucocorticoid Receptor alpha antibody - ChIP Grade (ab3580)

This image is courtesy of an anonymous Abreview

ab3580 staining Glucocorticoid Receptor alpha in mouse epididymis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with Bouin's solution and blocked with 1.5% serum for 30 minutes at 25°C; antigen retrieval was by heat mediation in a citrate buffer. Samples were incubated with primary antibody (1/1000 in blocking buffer) for 14 hours at 4°C. **ab6721** Goat **anti-rabbit HRP** (1/200) was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glucocorticoid Receptor alpha antibody - ChIP Grade (ab3580)

ab3580 (1µg/ml) staining glucocorticoid receptor alpha in human hippocampus using an automated system (DAKO Autostainer Plus). Using this protocol there is cytoplasmic staining in the neuropil and blood vessel smooth muscle. Sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffer EDTA pH 9.0 in a DAKO PT Link. Slides were peroxidase blocked in 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that, for manual staining, optimization of primary antibody concentration and incubation time is recommended. Signal amplification may be required.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors