

Anti-Glucocorticoid Receptor antibody [BuGR2] ab2768

★★★★★ [4 Abreviews](#) [46 References](#) [8 图像](#)

概述

产品名称	Anti-Glucocorticoid Receptor抗体[BuGR2]
描述	小鼠单克隆抗体[BuGR2] to Glucocorticoid Receptor
宿主	Mouse
特异性	Immunocytochemical staining of GR in L929 cells with this antibody results in staining of both the cytoplasm and nucleus, even in the presence of hormone. This antibody, using enzymatic digestion analysis, has been shown to react with the undigested 97 kDa GR, a 17 kDa DNA-binding trypsin fragment, and a 45 kDa steroid- and DNA-binding chymotrypsin fragment.
经测试应用	适用于: ICC/IF, Flow Cyt, WB
种属反应性	与反应: Mouse, Rat, Human 不与反应: Bird, Non human primates, Amphibian
免疫原	Full length protein corresponding to Rat Glucocorticoid Receptor. Partially purified rat GR.
阳性对照	ICC/IF: A549, HeLa, and U251 cells; Flow Cyt: Jurkat, HeLa, and NIH/3T3 cells; WB: Mouse liver lysate.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.05% Sodium azide Constituent: PBS
纯度	Protein A purified
克隆	单克隆

克隆编号	BuGR2
同种型	IgG2

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab2768于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF	★★★★★ (1)	1/50 - 1/500.
Flow Cyt		Use 0.5-1µg for 10 ⁶ cells. ab18414 - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
WB	★★★★☆ (2)	Use a concentration of 5 µg/ml. Detects a band of approximately 97 kDa (predicted molecular weight: 86 kDa). Using enzymatic digestion analysis detects a band of approximately 97 kDa, a 17 kDa DNA-binding trypsin fragment, and a 45 kDa steroid- and DNA-binding chymotrypsin fragment (predicted molecular weight: 86 kDa).

靶标


功能	<p>Receptor for glucocorticoids (GC). Has a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE) and as a modulator of other transcription factors. Affects inflammatory responses, cellular proliferation and differentiation in target tissues. Could act as a coactivator for STAT5-dependent transcription upon growth hormone (GH) stimulation and could reveal an essential role of hepatic GR in the control of body growth. Involved in chromatin remodeling. Plays a significant role in transactivation. Involved in nuclear translocation.</p>
组织特异性	<p>Widely expressed. In the heart, detected in left and right atria, left and right ventricles, aorta, apex, intraventricular septum, and atrioventricular node as well as whole adult and fetal heart.</p>
疾病相关	<p>Defects in NR3C1 are a cause of glucocorticoid resistance (GCRES) [MIM:138040]; also known as cortisol resistance. It is a hypertensive, hyperandrogenic disorder characterized by increased serum cortisol concentrations. Inheritance is autosomal dominant.</p>
序列相似性	<p>Belongs to the nuclear hormone receptor family. NR3 subfamily. Contains 1 nuclear receptor DNA-binding domain.</p>
结构域	<p>Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain.</p>
翻译后修饰	<p>Increased proteasome-mediated degradation in response to glucocorticoids. Phosphorylated in the absence of hormone; becomes hyperphosphorylated in the presence of glucocorticoid. The Ser-203-phosphorylated form is mainly cytoplasmic, and the Ser-211-phosphorylated form is nuclear. Transcriptional activity correlates with the amount of phosphorylation at Ser-211. Sumoylated; this reduces transcription transactivation. Ubiquitinated; restricts glucocorticoid-mediated transcriptional signaling.</p>

图片




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor antibody [BuGR2] - ChIP Grade (ab2768)

Immunocytochemistry/Immunofluorescence analysis of Glucocorticoid Receptor shows staining in A549 cells. Glucocorticoid Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with ab2768 (1:100) over night at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor antibody [BuGR2] - ChIP Grade (ab2768)

Immunocytochemistry/Immunofluorescence analysis of Glucocorticoid Receptor shows staining in HeLa cells. Glucocorticoid Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with ab2768 (1:100) over night at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.




Immunocytochemistry/ Immunofluorescence - Anti-Glucocorticoid Receptor antibody [BuGR2] - ChIP Grade (ab2768)

Immunocytochemistry/Immunofluorescence analysis of


Glucocorticoid Receptor shows staining in U251 cells.

Glucocorticoid Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with ab2768 (1:100) over night at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.




Flow Cytometry - Anti-Glucocorticoid Receptor antibody [BuGR2] - ChIP Grade (ab2768)

Flow cytometry analysis of Glucocorticoid Receptor showing positive staining in the nucleus and cytoplasm of NIH/3T3 cells compared to an isotype control (blue). Cells were harvested and adjusted to a concentration of $1-5 \times 10^6$ cells/ml. Cells were then fixed with 2% paraformaldehyde and washed with PBS. Cells were penetrated by dropping the supernatant and adding 90% methanol followed by incubation for 10 minutes at room temperature. Cells were blocked with a 2% solution of BSA-PBS for 30 min at room temperature and incubated with ab2768 at 2 µg/test for 60 min at room temperature. Cells were then incubated for 40 min at room temperature in the dark using a Dylight 488-conjugated goat anti-mouse IgG (H+L) secondary antibody and re-suspended in PBS for FACS analysis.



Flow Cytometry - Anti-Glucocorticoid Receptor antibody [BuGR2] - ChIP Grade (ab2768)


Flow cytometry analysis of Glucocorticoid Receptor showing positive staining in the nucleus and cytoplasm of Jurkat cells compared to an isotype control (blue). Cells were harvested and adjusted to a concentration of $1-5 \times 10^6$ cells/ml. Cells were then fixed with 2% paraformaldehyde and washed with PBS. Cells were penetrated by dropping the supernatant and adding 90% methanol followed by incubation for 10 minutes at room temperature. Cells were blocked with a 2% solution of BSA-PBS for 30 min at room temperature and then incubated with ab2768 at 1 µg/test for 60 min at room temperature. Cells were then incubated for 40 min at room temperature in the dark using a Dylight 488-conjugated goat anti-mouse IgG (H+L) secondary antibody and re-suspended in PBS for FACS analysis.



Cell: HeLa
Concentration: 1µg/test (100µl)
Theory location : Cytoplasm/Nucleus


Flow Cytometry - Anti-Glucocorticoid Receptor
antibody [BuGR2] - ChIP Grade (ab2768)

Flow cytometry analysis of Glucocorticoid Receptor showing positive staining in the nucleus and cytoplasm of HeLa cells compared to an isotype control (blue). Cells were harvested and adjusted to a concentration of $1-5 \times 10^6$ cells/ml. Cells were then fixed with 2% paraformaldehyde and washed with PBS. Cells were penetrated by dropping the supernatant and adding 90% methanol followed by incubation for 10 minutes at room temperature. Cells were blocked with a 2% solution of BSA-PBS for 30 min at room temperature and incubated with ab2768 at 1 µg/test for 60 min at room temperature. Cells were then incubated for 40 min at room temperature in the dark using a Dylight 488-conjugated goat anti-mouse IgG (H+L) secondary antibody and re-suspended in PBS for FACS analysis.



Western blot - Anti-Glucocorticoid Receptor
antibody [BuGR2] - ChIP Grade (ab2768)

Western blot of glucocorticoid receptor on mouse liver extract using ab2768. Western blot of glucocorticoid receptor on mouse liver extract using ab2768.



Flow Cytometry - Anti-Glucocorticoid Receptor
antibody [BuGR2] - ChIP Grade (ab2768)

Overlay histogram showing Jurkat cells stained with ab2768 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab2768, 0.5µg/ 1×10^6 cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2 (1µg/ 1×10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in Jurkat cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors