


### Anti-HDAC2 antibody [3F3] ab51832

敲除 验证

★★★★☆ 8 Abreviews 27 References 4 图像

#### 概述

产品名称	Anti-HDAC2抗体[3F3]
描述	小鼠单克隆抗体[3F3] to HDAC2
宿主	Mouse
经测试应用	适用于: Flow Cyt (Intra), WB, ICC/IF
种属反应性	与反应: Mouse, Human 预测可用于: Rat, Chicken, Cow 
免疫原	Synthetic peptide corresponding to Human HDAC2 aa 450-550.
阳性对照	Nuclear extract of HeLa cells and HAP1 whole cell lysate . IF/ICC: MCF7 cell line.
常规说明	<p>This monoclonal antibody to HDAC2 has been knockout validated in Western blot. The expected band for HDAC2 was observed in wild type cells and the band was not seen in HDAC2 knockout cells.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS

纯度	Protein G purified
克隆	单克隆
克隆编号	3F3
同种型	IgG1

## 应用

**The Abpromise guarantee**      **Abpromise™** 承诺保证使用 ab51832 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use 1µg for 10 <sup>6</sup> cells. <b>ab170190</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (4)	1/1000 - 1/10000. Detects a band of approximately 55 kDa (predicted molecular weight: 55 kDa).
ICC/IF	★★★★★ (1)	1/25 - 1/100.

## 靶标


<b>功能</b>	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1, PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity.
<b>组织特异性</b>	Widely expressed; lower levels in brain and lung.
<b>序列相似性</b>	Belongs to the histone deacetylase family. HD type 1 subfamily.
<b>翻译后修饰</b>	S-nitrosylated by GAPDH. In neurons, S-Nitrosylation at Cys-262 and Cys-274 does not affect the enzyme activity but abolishes chromatin-binding, leading to increases acetylation of histones and activate genes that are associated with neuronal development. In embryonic cortical neurons, S-Nitrosylation regulates dendritic growth and branching.
<b>细胞定位</b>	Nucleus.

## 图片



Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [3F3] (ab51832)

ab51832 staining HDAC2 in MCF7 cells. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab51832 at 5µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with **ab150117**, Goat polyclonal Secondary Antibody to Mouse IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour magenta). Nuclear DNA was labelled with DAPI (shown in blue). Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Western blot - Anti-HDAC2 antibody [3F3] (ab51832)

**Lane 1:** Wild type HAP1 whole cell lysate (20 µg)


**Lane 2:** HDAC2 knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** A431 whole cell lysate (20 µg)

**Lane 4:** HeLa whole cell lysate (20 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab51832 observed at 65 kDa. Red - loading control, **ab181602**, observed at 37 kDa.

ab51832 detected the expected band for HDAC2 in wild-type HAP1 cells and the band was not seen in HDAC2 knockout HAP1 cells. Wild-type and HDAC2 knockout samples were subjected to SDS-PAGE. ab51832 and **ab181602** (Rabbit anti GAPDH loading control) were incubated overnight at 4°C at 1000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed **ab216772** and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed **ab216777** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry (Intracellular) - Anti-HDAC2 antibody [3F3] (ab51832)

Overlay histogram showing HeLa cells stained with ab51832 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab51832, 1µg/1x10<sup>6</sup> cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (**ab91353**, 2µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed.



Western blot - Anti-HDAC2 antibody [3F3] (ab51832)

**All lanes** : Anti-HDAC2 antibody [3F3] (ab51832) at 5 µg/ml

**Lane 1** : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

**Lane 2** : HeLa (Human epithelial carcinoma cell line) Nuclear Lysate

**Lane 3** : Jurkat (Human T cell lymphoblast-like cell line) Whole Cell Lysate

**Lane 4** : NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

**Lane 5** : K562 (Human erythromyeloblastoid leukemia cell line) Whole Cell Lysate

**Lane 6** : U2OS (Human osteosarcoma cell line) Whole Cell Lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes** : Goat Anti-Mouse IgG H&L (HRP) preadsorbed (**ab97040**) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 55 kDa

**Observed band size:** 65 kDa

**Exposure time:** 16 minutes

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab51832 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors