

Anti-Hepatitis C Virus NS3 antibody ab21124

6 References

概述

产品名称	抗丙型肝炎病毒NS3抗体
描述	山羊多克隆抗体to丙型肝炎病毒NS3
宿主	Goat
经测试应用	适用于: ELISA, ICC/IF, WB
种属反应性	与反应: Hepatitis C virus
免疫原	Recombinant full length protein (genotype 1a).
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Preservative: 0.1% Sodium azide Constituent: 0.0268% PBS
纯度	Ion Exchange Chromatography
纯化说明	Sodium sulfate precipitation & ion-exchange chromatography.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab21124于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent dilution.
ICC/IF		Use at an assay dependent dilution.
WB		Use at an assay dependent dilution. Predicted molecular weight: 72 kDa.

靶标

相关性 HCV is a positive, single-stranded RNA virus in the Flaviviridae family. The genome is approximately 10,000 nucleotides and encodes a single polyprotein of about 3,000 amino acids. The polyprotein is processed by host cell and viral proteases into three major structural proteins including NS3, and several non-structural proteins necessary for viral replication. The NS3 part of the polyprotein displays three enzymatic activities: serine protease, NTPase and RNA helicase. The NS3 serine proteinase (NS3P) is a non-structural hepatitis C protein responsible for proteolytic processing of other non-structural proteins; because of this, it is also the most extensively studied protein of the Hepatitis C genome. It is responsible for proteolytic processing of the entire downstream region of the HC polyprotein, catalyzing cleavage at the NS3/NS4a, NS4a/NS4b, NS4b/NS5a, and NS5a/NS5b sites to release the mature NS3, NS4a, NS4b, NS5a, and NS5b proteins. For proper function, NS3 requires NS4a as a cofactor, but, interestingly enough, NS3 also cleaves the NS4a protein. The molecular weight of the monomer NS3P is 70 kDa.

细胞定位 Endoplasmic reticulum membrane

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors