

Anti-Histone H2A-Bbd antibody ab4175

3 References **2 图像**

概述

| | |
|-------|---|
| 产品名称 | Anti-Histone H2A-Bbd抗体 |
| 描述 | 兔多克隆抗体to Histone H2A-Bbd |
| 宿主 | Rabbit |
| 特异性 | Known to detect the recombinant form of human Histone H2A BBD. However, no reaction has yet been demonstrated against the endogenous protein in histone preparations. We believe that this is because of the low abundance of this variant in such preparations. |
| 经测试应用 | 适用于: ICC, WB |
| 种属反应性 | 与反应: Mouse, Chicken, Human |
| 免疫原 | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. |
| 常规说明 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

性能

| | |
|------|--|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |
| 存储溶液 | <p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: PBS</p> <p>Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.</p> |
| 纯度 | Immunogen affinity purified |
| 克隆 | 多克隆 |

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab4175于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|-----|------|---|
| ICC | | Use a concentration of 3 µg/ml. Staining may not be specific for H2A.Bbd (see image). |
| WB | | 1/500. Detects a band of approximately 12.5 kDa (predicted molecular weight: 12.5 kDa). As the WB shows, we can not find any reaction with any of the different histone extracts that we have used from different sources (calf, human, chicken). However, the antibody is very specific against the recombinant form of the protein. |

靶标

| | |
|-------|--|
| 功能 | Atypical histone H2A which can replace conventional H2A in some nucleosomes and is associated with active transcription and mRNA processing. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. Nucleosomes containing this histone are less rigid and organize less DNA than canonical nucleosomes in vivo. They are enriched in actively transcribed genes and associate with the elongating form of RNA polymerase. They associate with spliceosome components and are required for mRNA splicing. May participate in spermatogenesis. |
| 组织特异性 | Present in mature sperm. |
| 序列相似性 | Belongs to the histone H2A family. |
| 结构域 | The docking domain is responsible for the weaker heterodimerization with H2B. |
| 细胞定位 | Nucleus. Chromosome. Associated with the active X chromosome and with autosomes, while it is absent from the inactive X chromosome and excluded from Barr bodies. |

图片



Western blot - Anti-Histone H2A-Bbd antibody
(ab4175)


Juan Ausio

H2A-Bbd (ab4175)

CE= chicken erythrocyte (10 and 2 micrograms) , HE= HeLa cells (10 micrograms) and 2A*= recombinant H2ABbd (the numbers underneath the lanes are the number of nanograms loaded in each lane).

The primary antibody dilution was 1:500 and the secondary was 1:3000.

As the WB shows, we can not find any reaction with any of the different histone extracts that we have used from different sources (calf, human, chicken). However, the antibody is very specific against the recombinant form of the protein.



Immunocytochemistry - Anti-Histone H2A-Bbd
antibody (ab4175)

This image is courtesy of Petra Hajkova, University of Cambridge

Mouse embryonic fibroblasts (MEFs) were fixed in 4%PFA, permeabilised in PBS 1%BSA 0.1%TX-100 and incubated with ab4175 (anti-H2A.Bbd) antibody o/n. The antibody appears to show nucleolar staining. The antibody gave the same staining pattern in different cell types. The dilution used was 3ug/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors