


Anti-Histone H4 antibody - ChIP Grade ab7311

★★★★★ [12 Abreviews](#) [102 References](#) [5 图像](#)

概述

产品名称	Anti-Histone H4抗体- ChIP Grade
描述	兔多克隆抗体to Histone H4 - ChIP Grade
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, ChIP, WB
种属反应性	与反应: Mouse, Cow, Human, Saccharomyces cerevisiae 预测可用于: Mammals 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Calf Thymus Histone Preparation; Hela whole cell extract
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: PBS</p> <p>Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.</p>
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab7311于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (1)	Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
ICC/IF	★★★★★ (2)	1/200.
ChIP	★★★★★ (4)	Use 2 µg for 25 µg of chromatin.
WB	★★★★★ (5)	1/500 - 1/1000. Detects a band of approximately 14 kDa (predicted molecular weight: 11 kDa).


靶标

功能	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
序列相似性	Belongs to the histone H4 family.
翻译后修饰	Acetylation at Lys-6 (H4K5ac), Lys-9 (H4K8ac), Lys-13 (H4K12ac) and Lys-17 (H4K16ac) occurs in coding regions of the genome but not in heterochromatin. Citrullination at Arg-4 (H4R3ci) by PAD4 impairs methylation. Monomethylation and asymmetric dimethylation at Arg-4 (H4R3me1 and H4R3me2a, respectively) by PRMT1 favors acetylation at Lys-9 (H4K8ac) and Lys-13 (H4K12ac). Demethylation is performed by JMJD6. Symmetric dimethylation on Arg-4 (H4R3me2s) by the PRDM1/PRMT5 complex may play a crucial role in the germ-cell lineage. Monomethylated, dimethylated or trimethylated at Lys-21 (H4K20me1, H4K20me2, H4K20me3). Monomethylation is performed by SET8. Trimethylation is performed by SUV420H1 and SUV420H2 and induces gene silencing. Ubiquitinated by the CUL4-DDB-RBX1 complex in response to ultraviolet irradiation. This may weaken the interaction between histones and DNA and facilitate DNA accessibility to repair proteins. Monoubiquitinated at Lys-92 of histone H4 (H4K91ub1) in response to DNA damage. The exact role of H4K91ub1 in DNA damage response is still unclear but it may function as a licensing signal for additional histone H4 post-translational modifications such as H4 Lys-21 methylation (H4K20me). Sumoylated, which is associated with transcriptional repression.
细胞定位	Nucleus. Chromosome.


图片



Chromatin was prepared from Hela cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10min. The ChIP was performed with 25µg of chromatin, 2µg of ab7311 (blue), and 20µl of Protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach). Primers and probes are located in the first kb of the transcribed region.



IHC image of Histone H4 staining in human breast carcinoma FFPE section, performed on a Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab7311, 1µg/ml, for 8 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



All lanes : Anti-Histone H4 antibody - ChIP Grade (ab7311) at 1/500 dilution

Lane 1 : Calf thymus histone extract

Lane 2 : Calf thymus histone extract with Human Histone H4 peptide ([ab21044](#)) at 1 µg/ml

Lane 3 : Calf thymus histone extract with Histone H4 peptide - acetyl K16 ([ab22376](#)) at 1 µg/ml

Lane 4 : Calf thymus histone extract with Human Histone H4 (acetyl K20) peptide ([ab20632](#)) at 1 µg/ml


Lysates/proteins at 0.5 µg per lane.

Secondary

All lanes : Alexa-Fluor Goat anti-rabbit IgG at 1/10000 dilution

Predicted band size: 11 kDa


Observed band size: 14 kDa



SKN-SH cells were fixed in 4% paraformaldehyde for 10 mins, permeabilized in PBS-0.5% Triton X-100 for 5 mins and incubated for 30 minutes with ab7311 (1/200 dilution). The slides were rinsed once in PBS-Triton (0.1%), twice in PBS then incubated with the secondary antibody for 30 mins. The DNA is stained with DAPI (blue). Clear nuclear staining with ab7311 can be seen (green).

Immunocytochemistry/ Immunofluorescence - Anti-Histone H4 antibody - ChIP Grade (ab7311)

This image is courtesy of Darin McDonald



Anti-Histone H4 antibody - ChIP Grade (ab7311) at 1/500 dilution + Human Bone Whole Tissue Lysate at 25 µg with BSA, 1 Hour, 23°C at 5 %

Developed using the ECL technique.

Predicted band size: 11 kDa

Exposure time: 1 minute

Western blot - Anti-Histone H4 antibody - ChIP Grade (ab7311)

This image is courtesy of an Abreview submitted by Rodrigo Alves.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors