

Anti-HLA-C antibody [EPR26121-73] ab307361

重组 RabMAb

9 图像

概述

产品名称	Anti-HLA-C抗体[EPR26121-73]
描述	兔单克隆抗体[EPR26121-73] to HLA-C
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P 不适用于: Flow Cyt or ICC/IF
种属反应性	与反应: Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Jurkat transfected with scrambled siRNA control, Jurkat transfected with siRNA specifically targeting HLA-C whole cell lysate, A375, HeLa, Human lymphoma, HL-60, Human colon, and HEK-293T IHC-P: Human tonsil, Human NKT lymphoma and Human colon tissues. IP: Jurkat cell.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR26121-73

同种型IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab307361于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Predicted molecular weight: 41 kDa.
IP		1/30.
IHC-P		1/4000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

应用说明 Is unsuitable for Flow Cyt or ICC/IF.

靶标

图片



All lanes : Anti-HLA-C antibody [EPR26121-73] (ab307361) at 1/1000 dilution

Lane 1 : Jurkat (human T cell leukemia T lymphocyte) transfected with scrambled siRNA control whole cell lysate 20 µg

Lane 2 : Jurkat transfected with siRNA specifically targeting HLA-C whole cell lysate 20 µg

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (**ab97051**) at 1/100000 dilution


Predicted band size: 41 kDa

Observed band size: 41 kDa

Exposure time: 26 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST

Exposure time: 26 seconds



Western blot - Anti-HLA-C antibody [EPR26121-73]
(ab307361)

All lanes : Anti-HLA-C antibody [EPR26121-73] (ab307361) at 1/1000 dilution

Lane 1 : Jurkat (human T cell leukemia T lymphocyte) whole cell lysate 80 µg

Lane 2 : A375 (human malignant melanoma epithelial cell) whole cell lysate 80 µg

Lane 3 : HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate 80 µg

Lane 4 : Human lymphoma tissue lysate 80 µg

Lane 5 : HL-60 (human acute promyelocytic leukemia promyeloblast) whole cell lysate 80 µg

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/100000 dilution


Predicted band size: 41 kDa

Observed band size: 41 kDa

Exposure time: 15 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST

Exposure time: 15 seconds



Western blot - Anti-HLA-C antibody [EPR26121-73]
(ab307361)

Anti-CD5 antibody [EPR26532-12] - BSA and Azide free (Detector) ([ab307261](#)) at 1/1000 dilution + Human colon tissue lysate 40 µg

Secondary

Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution


Predicted band size: 41 kDa

Observed band size: 41 kDa

Exposure time: 7 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST

Exposure time: 7.25 seconds



All lanes : Anti-HLA-C antibody [EPR26121-73] (ab307361) at 1/1000 dilution

Lane 1 : HEK-293T cells transfected with an empty vector containing a his tag whole cell lysate 20 µg

Lane 2 : HEK-293T cells transfected with a human HLA-C expression vector containing a his tag whole cell lysate 20 µg

Lane 3 : HEK-293T cells transfected with a human HLA-A expression vector containing a his tag whole cell lysate 20 µg

Lane 4 : HEK-293T cells transfected with a human HLA-B expression vector containing a his tag whole cell lysate 20 µg

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/100000 dilution

Predicted band size: 41 kDa


Observed band size: 41 kDa

Exposure time: 3 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST

This antibody does not cross-react with human HLA-A and HLA-B.

Exposure time: 3 seconds




Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling HLA-C with ab307361 at 1/4000 (0.126 µg/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human tonsil (PMID: 17008885).

The section was incubated with ab307361 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 10 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HLA-C antibody [EPR26121-73] (ab307361)

Immunohistochemical analysis of paraffin-embedded Human NKT lymphoma tissue labeling HLA-C with ab307361 at 1/4000 (0.126 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on NKT lymphoma. The section was incubated with ab307361 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 10 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HLA-C antibody [EPR26121-73] (ab307361)

Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling HLA-C with ab307361 at 1/4000 (0.126 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human colon (PMID: 17008885). The section was incubated with ab307361 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 10 mins



Immunoprecipitation - Anti-HLA-C antibody
[EPR26121-73] (ab307361)

HLA-C was immunoprecipitated from 0.35 mg Jurkat (human T cell leukemia T lymphocyte) whole cell lysate 5 ug with ab307361 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab307361 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(**ab131366**) was used at 1/5000 dilution.

Lane 1: Jurkat (human T cell leukemia T lymphocyte) whole cell lysate 5 ug


Lane 2: abab307361 IP in Jurkat whole cell lysate

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab307361 in Jurkat whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds

Why choose a recombinant antibody?



Anti-HLA-C antibody [EPR26121-73] (ab307361)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors