

# Anti-HLA G antibody [EPR23298-97] ab283260

**重组** RabMAb

**2 References**   **7 图像**

### 概述

产品名称	Anti-HLA G抗体[EPR23298-97]
描述	兔单克隆抗体[EPR23298-97] to HLA G
宿主	Rabbit
经测试应用	<b>适用于:</b> IHC-P, WB, ICC/IF <b>不适用于:</b> Flow Cyt or IP
种属反应性	<b>与反应:</b> Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: JEG-3 whole cell lysate; Human placenta tissue lysate. IHC-P: Human placenta, Human breast carcinoma and Human cardiac muscle tissues. ICC/IF: JEG-3 cells.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <a href="#">see here</a> . Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb® patents</a> .

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59.94% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR23298-97

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab283260于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		1/20000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		1/1000. Predicted molecular weight: 38 kDa.
ICC/IF		1/100.


应用说明

Is unsuitable for Flow Cyt or IP.

靶标

功能	Involved in the presentation of foreign antigens to the immune system. Plays a role in maternal tolerance of the fetus by mediating protection from the deleterious effects of natural killer cells, cytotoxic T lymphocytes, macrophages and mononuclear cells.
组织特异性	Expressed in trophoblasts.
序列相似性	Belongs to the MHC class I family. Contains 1 Ig-like C1-type (immunoglobulin-like) domain.
细胞定位	Membrane.


图片



Immunohistochemical analysis of paraffin-embedded Human placenta tissue labelling HLA G with ab283260 at 1/20000 (0.026 µg/mL) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used. Positive staining on extravillous trophoblasts (EVTs) in human placenta (PMID:18538632, PMID:29249463, PMID:29249436). The section was incubated with ab283260 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Western blot - Anti-HLA G antibody [EPR23298-97] (ab283260)

**All lanes :** Anti-HLA G antibody [EPR23298-97] (ab283260) at 1/1000 dilution

**Lane 1 :** JEG-3 (human placenta choriocarcinoma epithelial cell) whole cell lysate

**Lane 2 :** HuT-78 (human Sezary syndrome cutaneous T lymphocyte) whole cell lysate

Lysates/proteins at 20 µg per lane.

## Secondary


**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

**Predicted band size:** 38 kDa

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Negative control: HuT-78 (PMID:2249951).


Exposure time: 48 seconds.



Immunocytochemistry/ Immunofluorescence - Anti-HLA G antibody [EPR23298-97] (ab283260)

Immunofluorescent analysis of 100% methanol-fixed, 0.1% TritonX-100 permeabilized JEG-3 cells labelling HLA G with ab283260 at 1/100 (5.18 µg/mL) dilution, followed by [ab150081](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 µg/mL) dilution (Green). Confocal image showing cytoplasmic and membranous staining in JEG-3 cell line. Negative control: HuT-78 (PMID: 2249951). [ab195889](#) Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5 µg/mL) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is [ab150081](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 µg/mL) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HLA G antibody [EPR23298-97] (ab283260)

Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue labelling HLA G with ab283260 at 1/20000 (0.026 µg/mL) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used. Positive staining on human breast carcinoma. The section was incubated with ab283260 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Western blot - Anti-HLA G antibody [EPR23298-97] (ab283260)

**All lanes :** Anti-HLA G antibody [EPR23298-97] (ab283260) at 1/1000 dilution

**Lane 1 :** Human placenta tissue lysate

**Lane 2 :** Human heart tissue lysate

Lysates/proteins at 20 µg per lane.

#### Secondary


**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

**Predicted band size:** 38 kDa

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Negative control: heart (PMID:22438923).

Exposure time: 26 seconds.







Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HLA G antibody [EPR23298-97] (ab283260)

Immunohistochemical analysis of paraffin-embedded Human cardiac muscle tissue labelling HLA G with ab283260 at 1/20000 (0.026 µg/mL) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used. Negative control: no staining on human cardiac muscle (PMID:22438923). The section was incubated with ab283260 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-HLA G antibody [EPR23298-97] (ab283260)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors