

# HRP Anti-Bid antibody [Y8] ab201754

**重组** RabMAb

## 1 References [3 图像](#)

### 概述

产品名称	HRP Anti-Bid抗体[Y8]
描述	HRP兔单克隆抗体[Y8] to Bid
宿主	Rabbit
偶联物	HRP
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Jurkat whole cell lysate. IHC-P: FFPE human colon adenocarcinoma tissue sections.
常规说明	Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> .

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C. Store In the Dark.
存储溶液	pH: 7.40 Preservative: 0.1% Proclin 300 Solution Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	Y8
同种型	IgG

### 应用

**The Abpromise guarantee**      **Abpromise<sup>™</sup>承诺保证使用ab201754于以下的经测试应用**


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/10000. Detects a band of approximately 22 kDa (predicted molecular weight: 22 kDa).
IHC-P		1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

## 靶标

<b>功能</b>	The major proteolytic product p15 BID allows the release of cytochrome c (By similarity). Isoform 1, isoform 2 and isoform 4 induce ICE-like proteases and apoptosis. Isoform 3 does not induce apoptosis. Counters the protective effect of Bcl-2.
<b>组织特异性</b>	Isoform 2 and isoform 3 are expressed in spleen, bone marrow, cerebral and cerebellar cortex. Isoform 2 is expressed in spleen, pancreas and placenta (at protein level). Isoform 3 is expressed in lung, pancreas and spleen (at protein level). Isoform 4 is expressed in lung and pancreas (at protein level).
<b>结构域</b>	Intact BH3 motif is required by BIK, BID, BAK, BAD and BAX for their pro-apoptotic activity and for their interaction with anti-apoptotic members of the Bcl-2 family.
<b>翻译后修饰</b>	TNF-alpha induces a caspase-mediated cleavage of p22 BID into a major p15 and minor p13 and p11 products. Phosphorylated upon DNA damage, probably by ATM or ATR. p15 BID is ubiquitinated by ITCH; ubiquitination results in proteasome-dependent degradation.
<b>细胞定位</b>	Cytoplasm; Cytoplasm. Mitochondrion membrane. When uncleaved, it is predominantly cytoplasmic; Mitochondrion membrane. A significant proportion of isoform 2 localizes to mitochondria, it may be cleaved constitutively; Mitochondrion membrane. Associated with the mitochondrial membrane and Mitochondrion membrane. Translocates to mitochondria as an integral membrane protein.

## 图片




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - HRP Anti-Bid antibody [Y8] (ab201754)

IHC image of Bid staining in a section of formalin-fixed paraffin-embedded colon adenocarcinoma\*, performed on a Leica BOND™. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab201754, 1/1000 dilution, for 15 mins at room temperature. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Western blot - HRP Anti-Bid antibody [Y8] (ab201754)

HRP Anti-Bid antibody [Y8] (ab201754) at 1/10000 dilution + Jurkat (Human T cell lymphoblast-like cell line) Whole Cell Lysate at 10 µg

Developed using the ECL technique.

Performed under reducing conditions.





**Predicted band size:** 22 kDa

**Observed band size:** 22 kDa

**Exposure time:** 4 seconds

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% milk before being incubated with ab201754 overnight at 4°C. Antibody binding was visualised using ECL development solution **ab133406**.

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

HRP Anti-Bid antibody [Y8] (ab201754)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors