


# HRP Anti-Histone H3 (di methyl K9) antibody [mAbcam 1220] ab62168

## 1 图像

### 概述

产品名称	HRP Anti-Histone H3 (di methyl K9)抗体[mAbcam 1220]
描述	HRP小鼠单克隆抗体[mAbcam 1220] to Histone H3 (di methyl K9)
宿主	Mouse
偶联物	HRP
经测试应用	适用于: WB
种属反应性	与反应: Cow 预测可用于: Mouse, Rat, Human, Saccharomyces cerevisiae, Xenopus laevis, Arabidopsis thaliana, Caenorhabditis elegans, Drosophila melanogaster, Schizosaccharomyces pombe, Mammals, Rice 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as <a href="#">ab1772</a> )
阳性对照	This antibody gave a positive signal in calf thymus histone preparation (Nuclear Lysate).
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Store In the Dark.
存储溶液	pH: 7.40 Preservative: 0.1% Proclin 300 Solution Constituents: PBS, 1% BSA, 30% Glycerol (glycerin, glycerine)
纯度	IgG fraction

克隆	单克隆
克隆编号	mAbcam 1220
骨髓瘤	Sp2/0
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab62168于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/20000 - 1/50000. Detects a band of approximately 17 kDa (predicted molecular weight: 15 kDa).Can be blocked with <b><u>Human Histone H3 (di methyl K9) peptide (ab1772)</u></b> .

靶标

功能	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
序列相似性	Belongs to the histone H3 family.
发展阶段	Expressed during S phase, then expression strongly decreases as cell division slows down during the process of differentiation.
翻译后修饰	<p>Acetylation is generally linked to gene activation. Acetylation on Lys-10 (H3K9ac) impairs methylation at Arg-9 (H3R8me2s). Acetylation on Lys-19 (H3K18ac) and Lys-24 (H3K24ac) favors methylation at Arg-18 (H3R17me).</p> <p>Citrullination at Arg-9 (H3R8ci) and/or Arg-18 (H3R17ci) by PAD4 impairs methylation and represses transcription.</p> <p>Asymmetric dimethylation at Arg-18 (H3R17me2a) by CARM1 is linked to gene activation. Symmetric dimethylation at Arg-9 (H3R8me2s) by PRMT5 is linked to gene repression. Asymmetric dimethylation at Arg-3 (H3R2me2a) by PRMT6 is linked to gene repression and is mutually exclusive with H3 Lys-5 methylation (H3K4me2 and H3K4me3). H3R2me2a is present at the 3' of genes regardless of their transcription state and is enriched on inactive promoters, while it is absent on active promoters.</p> <p>Methylation at Lys-5 (H3K4me), Lys-37 (H3K36me) and Lys-80 (H3K79me) are linked to gene activation. Methylation at Lys-5 (H3K4me) facilitates subsequent acetylation of H3 and H4. Methylation at Lys-80 (H3K79me) is associated with DNA double-strand break (DSB) responses and is a specific target for TP53BP1. Methylation at Lys-10 (H3K9me) and Lys-28 (H3K27me) are linked to gene repression. Methylation at Lys-10 (H3K9me) is a specific target for HP1 proteins (CBX1, CBX3 and CBX5) and prevents subsequent phosphorylation at Ser-11 (H3S10ph) and acetylation of H3 and H4. Methylation at Lys-5 (H3K4me) and Lys-80 (H3K79me) require preliminary monoubiquitination of H2B at 'Lys-120'. Methylation at Lys-10 (H3K9me) and Lys-28 (H3K27me) are enriched in inactive X chromosome chromatin.</p> <p>Phosphorylated at Thr-4 (H3T3ph) by GSG2/haspin during prophase and dephosphorylated</p>


during anaphase. Phosphorylation at Ser-11 (H3S10ph) by AURKB is crucial for chromosome condensation and cell-cycle progression during mitosis and meiosis. In addition phosphorylation at Ser-11 (H3S10ph) by RPS6KA4 and RPS6KA5 is important during interphase because it enables the transcription of genes following external stimulation, like mitogens, stress, growth factors or UV irradiation and result in the activation of genes, such as c-fos and c-jun. Phosphorylation at Ser-11 (H3S10ph), which is linked to gene activation, prevents methylation at Lys-10 (H3K9me) but facilitates acetylation of H3 and H4. Phosphorylation at Ser-11 (H3S10ph) by AURKB mediates the dissociation of HP1 proteins (CBX1, CBX3 and CBX5) from heterochromatin. Phosphorylation at Ser-11 (H3S10ph) is also an essential regulatory mechanism for neoplastic cell transformation. Phosphorylated at Ser-29 (H3S28ph) by MLTK isoform 1, RPS6KA5 or AURKB during mitosis or upon ultraviolet B irradiation. Phosphorylation at Thr-7 (H3T6ph) by PRKCBB is a specific tag for epigenetic transcriptional activation that prevents demethylation of Lys-5 (H3K4me) by LSD1/KDM1A. At centromeres, specifically phosphorylated at Thr-12 (H3T11ph) from prophase to early anaphase, by DAPK3 and PKN1. Phosphorylation at Thr-12 (H3T11ph) by PKN1 is a specific tag for epigenetic transcriptional activation that promotes demethylation of Lys-10 (H3K9me) by KDM4C/JMJD2C. Phosphorylation at Tyr-42 (H3Y41ph) by JAK2 promotes exclusion of CBX5 (HP1 alpha) from chromatin.

Monoubiquitinated by RAG1 in lymphoid cells, monoubiquitination is required for V(D)J recombination (By similarity). Ubiquitinated by the CUL4-DDB-RBX1 complex in response to ultraviolet irradiation. This may weaken the interaction between histones and DNA and facilitate DNA accessibility to repair proteins.

## 细胞定位

Nucleus. Chromosome.

## 图片



**All lanes :** HRP Anti-Histone H3 (di methyl K9) antibody [mAbcam 1220] (ab62168) at 1/20000 dilution

**Lane 1 :** Calf Thymus Histone Preparation Nuclear Lysate

**Lane 2 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (unmodified ) peptide ([ab2903](#)) at 2.5 µg/ml

**Lane 3 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (mono methyl K4) peptide ([ab1340](#)) at 2.5 µg/ml

**Lane 4 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (di methyl K4) peptide ([ab7768](#)) at 2.5 µg/ml

**Lane 5 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (tri methyl K4) peptide ([ab1342](#)) at 2.5 µg/ml

**Lane 6 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (mono methyl K9) peptide ([ab1771](#)) at 2.5 µg/ml

**Lane 7 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (di methyl K9) peptide ([ab1772](#)) at 2.5 µg/ml

**Lane 8 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (tri methyl K9) peptide ([ab1773](#)) at 2.5 µg/ml

**Lane 9 :** Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (mono methyl K27) peptide ([ab1780](#)) at 2.5

µg/ml

**Lane 10** : Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (di methyl K27) peptide (**ab1781**) at 2.5 µg/ml

**Lane 11** : Calf Thymus Histone Preparation Nuclear Lysate with Human Histone H3 (tri methyl K27) peptide (**ab1782**) at 2.5 µg/ml

Lysates/proteins at 0.5 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 15 kDa

**Observed band size:** 17 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors