abcam

Product datasheet

HRP Anti-Iba1 antibody [EPR16589] ab216544



重组 RabMAb

2 图像

概述

产品名称 HRP Anti-lba1抗体[EPR16589]

描述 HRP兔单克隆抗体[EPR16589] to lba1

宿主 Rabbit 偶联物 HRP

经测试应用 适用于: IHC-P 种属反应性

与反应: Human

预测可用于: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 IHC-P: normal human hippocampus tissue sections

This product is a recombinant monoclonal antibody, which offers several advantages including: 常规说明

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information **see here**.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式 Liquid

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. 存放说明

Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 1% BSA, 30% Glycerol (glycerin, glycerine), PBS

纯度 Protein A purified

克隆 单克隆

克隆编号 EPR16589

同种型 lgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab216544于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
IHC-P		1/50. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

***	_		_
31	ш	*	

功能	Actin-binding protein that enhances membrane ruffling and RAC activation. Enhances the actin-bundling activity of LCP1. Binds calcium. Plays a role in RAC signaling and in phagocytosis. May play a role in macrophage activation and function. Promotes the proliferation of vascular smooth muscle cells and of T-lymphocytes. Enhances lymphocyte migration. Plays a role in vascular inflammation.
组织 特异性	Detected in T-lymphocytes and peripheral blood mononuclear cells.
序列相似性	Contains 2 EF-hand domains.
翻译后修饰	Phosphorylated on serine residues.
细胞定位	Cytoplasm > cytoskeleton. Cell projection > ruffle membrane. Associated with the actin

cytoskeleton at membrane ruffles and at sites of phagocytosis.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - HRP Anti-lba1 antibody
[EPR16589] (ab216544)

IHC image of lba1 staining in a section of formalin-fixed paraffinembedded normal human hippocampus, performed on a Leica BOND™. The section was pre-treated using heat mediated antigen retrieval with Tris/EDTA buffer (pH9, epitope retrieval solution 2) for 20mins. The section was then incubated with ab216544, 1/50 dilution, for 15 mins at room temperature. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors