abcam

Product datasheet

HRP Anti-PDGFR alpha + PDGFR beta antibody [Y92] ab195515



重组 RabMAb

2 图像

概述

产品名称 HRP Anti-PDGFR alpha + PDGFR beta抗体[Y92]

描述 HRP兔单克隆抗体[Y92] to PDGFR alpha + PDGFR beta

宿主 Rabbit 偶联物 HRP

经测试应用 适用干: IHC-P 种属反应性 与反应: Human

预测可用于: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 IHC-P: FFPE human normal kidney tissue sections. WB: N-GST tagged Human PDGF Receptor

beta (aa557 to 1106) recombinant protein, N-GST tagged Human PDGF Receptor alpha (aa550

to 1089) recombinant protein.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity - Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

纯**度** Protein A purified

克隆 单克隆

克隆编号 Y92

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab195515于以下的经测试应用


"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		1/100. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

靶标

细胞定位 PDGFR alpha: Membrane. PDGFR beta: Membrane.

图片




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - HRP Anti-PDGFR alpha + PDGFR beta antibody [Y92] (ab195515)

IHC image of PDGFR alpha + beta staining in a section of formalin-fixed paraffin-embedded human normal kidney*. The section was pre-treated using pressure cooker heat mediated antigen retrieval with sodium citrate buffer (pH6) for 30mins, and incubated overnight at +4°C with ab195515 at 1µg/ml. DAB was used as the chromogen (ab103723), diluted 1/100 and incubated for 10min at room temperature. The section was counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors