


Anti-IL-21 antibody ab5978

★★★★★ [2 Abreviews](#) [9 References](#) [4 图像](#)

概述

产品名称	Anti-IL-21抗体
描述	兔多克隆抗体to IL-21
宿主	Rabbit
经测试应用	适用于: ICC/IF, WB
种属反应性	与反应: Human 预测可用于: Cow, Dog, Pig 
免疫原	Synthetic peptide corresponding to Human IL-21 aa 121-135. Sequence: TCPSCDSYEKKPPKE

Database link: [Q9HBE4](#)

(Peptide available as [ab6025](#), [ab6680](#))

 [Run BLAST with](#)

 [Run BLAST with](#)

常规说明

A novel cytokine was recently identified in human and mouse and designated IL-21, which has significant homology to IL-2, IL-4, and IL-15. The receptor for IL-21 (IL-21R, also termed NLR for novel Interleukin receptor) is a new member of the class I cytokine receptor family. IL-21R forms a complex with the common cytokine receptor g chain, gc, and mediates IL-21 signaling. IL-21 and its receptor activate JAK-STAT signaling pathway. IL-21 is expressed in activated T cells, and HL-60 and THP-1 cell lines. IL-21 plays a role in the proliferation and maturation of NK, B and T cell populations.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C long term. Avoid freeze /

	thaw cycle.
存储溶液	pH: 7.2 Preservative: 0.02% Sodium azide
纯度	Immunogen affinity purified
Primary antibody说明	A novel cytokine was recently identified in human and mouse and designated IL-21, which has significant homology to IL-2, IL-4, and IL-15. The receptor for IL-21 (IL-21R, also termed NLR for novel Interleukin receptor) is a new member of the class I cytokine receptor family. IL-21R forms a complex with the common cytokine receptor g chain, gc, and mediates IL-21 signaling. IL-21 and its receptor activate JAK-STAT signaling pathway. IL-21 is expressed in activated T cells, and HL-60 and THP-1 cell lines. IL-21 plays a role in the proliferation and maturation of NK, B and T cell populations.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab5978于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use a concentration of 2 µg/ml.
WB	★★★★★ (1)	Use a concentration of 0.5 - 1 µg/ml. Detects a band of approximately 18 kDa.

靶标

功能	Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL 15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL 15 and IL 18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation.
组织特异性	Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes.
序列相似性	Belongs to the IL-15/IL-21 family.
细胞定位	Secreted.


图片




All lanes : Anti-IL-21 antibody (ab5978) at 1 µg/mL, 1h incubation at RT in 5% NFM/TBST.

Loading: 15 µg of lysates per lane.

Secondary: Goat anti-rabbit IgG HRP conjugate at 1/10000 dilution.



Western blot analysis of IL-21 expression in HL-60 cell lysate in the absence (lane A) or presence of blocking peptide (lane B) with ab5978 at 1 µg /ml. Western blot analysis of IL-21 expression in HL-60 cell lysate in the absence (lane A) or presence of blocking peptide (lane B) with ab5978 at 1 µg /ml.



Anti-IL-21 antibody (ab5978) at 1 µg/ml + Recombinant human IL-21 protein ([ab73250](#)) at 0.1 µg


Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ([ab97080](#)) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Exposure time: 8 minutes



ab5978 at 2ug/ml staining of IL21 in K562 cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors