

### Anti-Leu Enkephalin antibody [NOC1] ab150346

★★★★★ [3 Abreviews](#) [9 References](#)

#### 概述

产品名称	Anti-Leu Enkephalin抗体[NOC1]
描述	小鼠单克隆抗体[NOC1] to Leu Enkephalin
宿主	Mouse
特异性	This antibody does not distinguish between Met-enkephalin and Leu-enkephalin by radioimmunoassay or by immunocytochemistry.
经测试应用	适用于: IHC-Fr, ICC, RIA, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide corresponding to Leu Enkephalin conjugated to bovine serum albumin. <a href="#">Run BLAST with ExPASy</a> <a href="#">Run BLAST with NCBI</a>
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Contains no additives.
纯度	Tissue culture supernatant
纯化说明	ab150346 is in a protein-free culture medium.
克隆	单克隆
克隆编号	NOC1
同种型	IgG1

## 应用

### The Abpromise guarantee

**Abpromise™**承诺保证使用ab150346于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.
RIA		Use at an assay dependent concentration.
IHC-P	★★★★★ (1)	Use at an assay dependent concentration.

## 靶标

相关性	Enkephalin and enkephalin-like peptides have been detected in many animal species. In mammals these opioid pentapeptides are cleaved from a common precursor molecule which is expressed in a number of cell types, including neurons and endocrine cells associated with the sympathetic nervous system where they are often co-stored with catecholamines. They are known to be endogenous ligands for Sopioid receptors and are known to mediate analgesia at the spinal level.
细胞定位	Secreted

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors