abcam

Product datasheet

Anti-LSP1 antibody [EPR5997] ab133506



★★★★★ 1 Abreviews 2 References 12 图像

概述

产品名称 Anti-LSP1抗体[EPR5997]

描述 兔单克隆抗体[EPR5997] to LSP1

宿主 Rabbit

经测试应用 适用于: WB, IHC-P, ICC/IF, Flow Cyt (Intra)

不适用干: IP

种属反应性 与反应: Mouse. Rat. Human

免疫原 Synthetic peptide within Human LSP1 aa 300-400 (C terminal). The exact sequence is

proprietary.

阳性对照 WB: Daudi, Ramos and HuT78 cell lysates. Mouse spleen and Rat brain tissue lysates. IHC-P:

Human colon, Mouse spleen, Rat spleen and Human tonsil tissue. ICC/IF: Daudi Flow Cyt (intra):

Daudi

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.


Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

性能

形式 Liquid

Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C. 存放说明

 $K_D = 1.02 \times 10^{-10} M$ 解离常数(KD)



Learn more about K_D

存储溶液 pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)

纯**度** Protein A purified

 克隆
 单克隆

 克隆编号
 EPR5997

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab133506于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/10000 - 1/50000. Detects a band of approximately 52 kDa (predicted molecular weight: 37 kDa).
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols.
ICC/IF		1/100. For unpurified use at 1/500 - 1/1000.
Flow Cyt (Intra)		1/100.

应**用说明** Is unsuitable for IP.

靶标

功能 May play a role in mediating neutrophil activation and chemotaxis.

组织**特异性** Activated T-lymphocytes.


翻译后修饰 Phosphorylated by casein kinase II, protein kinase C and MAPKAPK2. Phosphorylation by PKC

induces translocation from membrane to cytoplasm. Phosphorylation by MAPKAPK2 may

regulate neutrophil chemotaxis.


细**胞定位** Cell membrane.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-LSP1 antibody
[EPR5997] (ab133506)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human colon tissue sections labeling LSP1 with purified ab133506 at 1:500 dilution (1.954 µg/ml). Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Western blot - Anti-LSP1 antibody [EPR5997] (ab133506)

Anti-LSP1 antibody [EPR5997] (ab133506) at 1/10000 dilution (Purified) + Rat brain lysates at 15 μg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 37 kDa



Western blot - Anti-LSP1 antibody [EPR5997] (ab133506)

All lanes : Anti-LSP1 antibody [EPR5997] (ab133506) at 1/10000 dilution (Purified)

Lane 1: Mouse spleen lysates

Lane 2 : Daudi (Human Burkitt's lymphoma lymphoblast) whole cell lysate


Lane 3: HuT-78 (Human Sezary syndrome cutaneous T lymphocyte) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit lgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution


Predicted band size: 37 kDa **Observed band size:** 52 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-LSP1 antibody

[EPR5997] (ab133506)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat spleen tissue sections labeling LSP1 with purified ab133506 at 1:500 dilution (1.954 μ g/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-LSP1 antibody


[EPR5997] (ab133506)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse spleen tissue sections labeling LSP1 with purified ab133506 at 1:500 dilution (1.954 µg/ml). Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunocytochemistry/ Immunofluorescence - Anti-LSP1 antibody [EPR5997] (ab133506)

Immunocytochemistry/ Immunofluorescence analysis of Daudi (Human Burkitt's lymphoma lymphoblast) cells labeling LSP1 with purified ab133506 at 1:100 dilution (9.8 μ g/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 μ g/ml). Goat anti rabbit lgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 μ g/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-LSP1 antibody [EPR5997] (ab133506)

Intracellular Flow Cytometry analysis of Daudi (Human Burkitt's lymphoma lymphoblast) cells labeling LSP1 with purified ab133506 at 1/100 dilution (10 μ g/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit lgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal lgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-LSP1 antibody [EPR5997] (ab133506)

All lanes : Anti-LSP1 antibody [EPR5997] (ab133506) at 1/10000 dilution


Lane 1 : Daudi cell lysate
Lane 2 : Ramos cell lysate
Lane 3 : HuT78 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: Goat-anti-rabbit HRP at 1/2000 dilution

Predicted band size: 37 kDa Observed band size: 52 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-LSP1 antibody

[EPR5997] (ab133506)

Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labelling LSP1 with ab133506 at 1/250 dilution.


Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



OI-RD Scanning - Anti-LSP1 antibody [EPR5997] (ab133506)

Equilibrium disassociation constant (K_D) Learn more about K_D


Click here to learn more about K_D



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-LSP1 antibody
[EPR5997] (ab133506)

Immunohistochemical analysis of paraffin-embedded Human colon tissue labelling LSP1 with ab133506 at 1/250 dilution.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors