

Anti-Matrin 3 antibody [EPR10635(B)] ab151714






[5 Abreviews](#)
[10 References](#)
[15 图像](#)

概述

产品名称	Anti-Matrin 3抗体[EPR10635(B)]
描述	兔单克隆抗体[EPR10635(B)] to Matrin 3
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	HEK293T, K562, HeLa and HepG2 whole cell lysate (ab7900); Human brain tissue, Human colon tissue and Human breast carcinoma tissue; Mouse and rat liver tissue, Mouse and Rat brain tissue lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR10635(B)
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab151714于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/220.
WB	★★★★★ (2)	1/10000 - 1/50000. Predicted molecular weight: 95 kDa.
IHC-P		1/2000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/250 - 1/500.
ICC/IF	★★★★★ (1)	1/250 - 1/500.

靶标

功能

May play a role in transcription or may interact with other nuclear matrix proteins to form the internal fibrogranular network. In association with the SFPQ-NONO heteromer may play a role in nuclear retention of defective RNAs.

疾病相关

Defects in MATR3 are the cause of myopathy distal type 2 (MPD2) [MIM:606070]; also called vocal cord and pharyngeal dysfunction with distal myopathy (VCPDM). MPD2 is a muscular disorder characterized by distal weakness, with onset in hands and feet, associated with vocal cord and pharyngeal weakness causing a nasal voice and swallowing disorders.


序列相似性

Contains 1 matrin-type zinc finger.
Contains 2 RRM (RNA recognition motif) domains.

细胞定位


Nucleus matrix.

图片



Western blot - Anti-Matrin 3 antibody [EPR10635(B)]
(ab151714)

ab151714 was shown to react with MATR3 in wild-type HAP1 cells in Western blot with loss of signal observed in a MATR3 knockout cell line. Wild-type HAP1 and MATR3 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab151714 overnight at 4 °C at a 1/10000 dilution. Blots were incubated with goat anti-rabbit HRP secondary antibodies at 0.2 µg/ml before imaging. These data were provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.



Western blot - Anti-Matrin 3 antibody [EPR10635(B)]
(ab151714)

All lanes : Anti-Matrin 3 antibody [EPR10635(B)] (ab151714) at 1/10000 dilution (purified)

Lane 1 : HEK-293 (Human epithelial cell line from embryonic kidney) whole cell lysate

Lane 2 : K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate

Lane 3 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 4 : Mouse brain tissue lysate

Lane 5 : Rat brain tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 95 kDa


Observed band size: 125 kDa

Blocking/Diluting buffer 5% NFDM/TBST




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Immunohistochemical analysis of paraffin-embedded human colon tissue sections labeling Matrin 3 with purified ab151714 at a dilution of 1/2000 (0.7 µg/ml). **ab97051** Goat Anti-Rabbit IgG H&L (HRP) at 1/500 was used as the secondary antibody. Sections were counterstained with hematoxylin. Antigen retrieval was heat mediated using citrate Buffer, pH 6.0. PBS was used instead of the primary antibody as the negative control and is shown in the inset.



Immunocytochemistry/ Immunofluorescence - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Immunocytochemistry/Immunofluorescence analysis of HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling Matrin 3 with purified ab151714 at 1/500. Cells were fixed with 4% Paraformaldehyde and permeabilised with 0.1% tritonX-100. **ab150077**, Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. Cells were counterstained with **ab195889** Anti-Alpha Tubulin antibody [DM1A] (1/200, 2.5 µg/mL) - Microtubule Marker (Alexa Fluor® 594) at 1/200. DAPI (blue) was used as a nuclear counterstain. Secondary Only Control: PBS was used instead of the primary antibody as the negative control.



All lanes : Anti-Matrin 3 antibody [EPR10635(B)] (ab151714) at 1/1000 dilution (unpurified)

Lane 1 : HEK293T cell lysate

Lane 2 : K562 cell lysate

Lane 3 : HeLa cell lysate


Lane 4 : HepG2 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 95 kDa




Immunohistochemical analysis of Paraffin-embedded rat liver tissue sections labeling Matrin 3 with purified ab151714 at a dilution of 1/2000 (0.7 µg/ml). **ab97051** Goat Anti-Rabbit IgG H&L (HRP) at 1/500 was used as the secondary antibody. Sections were counterstained with hematoxylin. Antigen retrieval was heat mediated using citrate Buffer, pH 6.0. PBS was used instead of the primary antibody as the negative control and is shown in the inset.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)



Immunohistochemical analysis of paraffin-embedded mouse liver tissue sections labeling Matrin 3 with purified ab151714 at a dilution of 1/2000 (0.7 µg/ml). **ab97051** Goat Anti-Rabbit IgG H&L (HRP) at 1/500 was used as the secondary antibody. Sections were counterstained with hematoxylin. Antigen retrieval was heat mediated using citrate Buffer, pH 6.0. PBS was used instead of the primary antibody as the negative control and is shown in the inset.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)



Intracellular Flow Cytometry analysis of 4%paraformaldehyde fixedHepG2 (human hepatocellular carcinoma) cells labeling Matrin 3 with purified ab151714 at a dilution of 1/220. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isootype control: Rabbit monoclonal IgG (Black)Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)


Flow Cytometry (Intracellular) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Immunohistochemical analysis of paraffin-embedded Human brain tissue labeling Matrin 3 with unpurified ab151714 at 1/250 dilution.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue labeling Matrin 3 with unpurified ab151714 at 1/250 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)


Immunofluorescent analysis of HepG2 cells labeling Matrin 3 with unpurified ab151714 at 1/250 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Immunohistochemical analysis of paraffin embedded normal Human uterus tissue using unpurified ab151714 showing +ve staining.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Immunohistochemical analysis of paraffin embedded Human Ovarian carcinoma tissue using unpurified ab151714 showing +ve staining.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemical analysis of paraffin embedded Human Lung carcinoma tissue using unpurified ab151714 showing +ve staining.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Why choose a recombinant antibody?



Anti-Matrin 3 antibody [EPR10635(B)] (ab151714)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors