


Anti-mH2A1 antibody [EPR9359(2)] ab183041

敲除验证
重组
RabMAb

★★★★★
[2 Abreviews](#)
[13 References](#)
[10 图像](#)

概述

产品名称	Anti-mH2A1 抗体[EPR9359(2)]
描述	兔单克隆抗体[EPR9359(2)] to mH2A1
宿主	Rabbit
经测试应用	适用于: WB, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Human 预测可用于: Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HepG2, MCF7, 293T and HeLa whole cell lysate (ab150035) IHC-P: Human kidney and liver tissues ICC-IF: HAP1-WT and H2AFY knockout cells. MCF7 and HeLa cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR9359(2)

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab183041于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/10000 - 1/50000. Detects a band of approximately 40 kDa (predicted molecular weight: 40 kDa).
IHC-P	★★★★★ (1)	1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF	★★★★★ (1)	Use a concentration of 1 µg/ml. This antibody gives positive signal in both 4%PFA and 100% MeOH-fixed cells.

靶标

功能	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes where it represses transcription. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Involved in stable X chromosome inactivation. Inhibits the binding of transcription factors and interferes with the activity of remodeling SWI/SNF complexes. Inhibits histone acetylation by EP300 and recruits class I HDACs, which induces an hypoacetylated state of chromatin. In addition, isoform 1, but not isoform 2, binds ADP-ribose and O-acetyl-ADP-ribose, and may be involved in ADP-ribose-mediated chromatin modulation.
组织特异性	Ubiquitous.
序列相似性	Contains 1 histone H2A domain. Contains 1 Macro domain.
翻译后修饰	Monoubiquitinated at either Lys-116 or Lys-117. May also be polyubiquitinated. Ubiquitination is mediated by the CUL3/SPOP E3 complex and does not promote proteasomal degradation. Instead, it is required for enrichment in inactive X chromosome chromatin.
细胞定位	Nucleus. Chromosome. Enriched in inactive X chromosome chromatin and in senescence-associated heterochromatin.

图片



Western blot - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

Lane 1: Wild type HAP1 whole cell lysate (20 µg)


Lane 2: mH2A1 knockout HAP1 whole cell lysate (20 µg)

Lane 3: HeLa whole cell lysate (20 µg)

Lane 4: Hepg2 whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab183041 observed at 40 kDa. Red - loading control, **ab18058**, observed at 130 kDa.


ab183041 was shown to specifically react with mH2A1 when mH2A1 knockout samples were used. Wild-type and mH2A1 knockout samples were subjected to SDS-PAGE. Ab183041 and **ab18058** (Mouse anti Vinculin loading control) were incubated overnight at 4°C at 10000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)


ab183041 staining mH2A1 in HAP1 WT and H2AFY knockout cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1%PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab183041 at 1µg/ml and **ab195889**, Mouse monoclonal [DM1A] to alpha Tubulin - Microtubule Marker (Alexa Fluor® 594) at 1/250 dilution (shown in pseudocolor red). Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488) at 1/1000 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling mH2A1 with ab183041 at 1/100 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.



Western blot - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

All lanes : Anti-mH2A1 antibody [EPR9359(2)] (ab183041) at 1/1000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : H2AFY CRISPR/Cas9 edited HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 40 kDa

Observed band size: 40 kDa

Lanes 1- 2: Merged signal (red and green). Green - ab183041 observed at 40 kDa. Red - Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) observed at 50 kDa.

ab183041 was shown to react with mH2A1 in wild-type HEK-293T cells in western blot. The band observed in CRISPR/Cas9 edited cell line [ab266241](#) (CRISPR/Cas9 edited cell lysate [ab257463](#)) lane below 40kDa may represent truncated forms and

cleaved fragments. This has not been investigated further. Wild-type HEK-293T and H2AFY CRISPR/Cas9 edited HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab183041 and Anti-alpha Tubulin antibody [DM1A] - Loading Control ([ab7291](#)) were incubated overnight at 4°C at a 1 in 1000 Dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



All lanes : Anti-mH2A1 antibody [EPR9359(2)] (ab183041) at 1/50000 dilution

Lane 1 : HepG2 cell lysate

Lane 2 : MCF7 cell lysate

Lane 3 : HeLa cell lysate

Lane 4 : 293T cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution


Predicted band size: 40 kDa

Observed band size: 40 kDa




Immunocytochemistry/ Immunofluorescence - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

Immunofluorescent analysis of 4% paraformaldehyde-fixed MCF7 cells labeling mH2A1 with ab183041 at 1/250 dilution followed by Goat anti rabbit IgG (Alexa Fluor® 488) at 1/200 dilution (green). Counter stained with Dapi (blue).




Immunocytochemistry/ Immunofluorescence - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

Immunofluorescent analysis of acetone-fixed HeLa cells labeling mH2A1 with ab183041 at 1/250 dilution followed by Goat anti rabbit IgG (Alexa Fluor® 488) at 1/200 dilution (green). Counter stained with Dapi (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling mH2A1 with ab183041 at 1/100 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

This image is courtesy of an anonymous abreview.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver from wild-type and mH2A1/2 knock out tissue sections labeling mH2A1 with ab183041 at 1/400 dilution. Sections were fixed in formaldehyde; heat mediated antigen retrieval was performed using a citrate buffer pH 6. An undiluted polyclonal horse anti-rabbit IgG (HRP-conjugated) was used as the secondary antibody.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-mH2A1 antibody [EPR9359(2)] (ab183041)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors