

### Anti-mTOR antibody [Y391] - BSA and Azide free ab218525

重组 RabMAb

#### 8 图像

#### 概述

产品名称	Anti-mTOR抗体[Y391] - BSA and Azide free
描述	兔单克隆抗体[Y391] to mTOR - BSA and Azide free
宿主	Rabbit
特异性	Expression levels of the target protein vary with sample type and some optimisation may be required.
经测试应用	<b>适用于:</b> IHC-Fr, WB, IP, IHC-P <b>不适用于:</b> Flow Cyt or ICC/IF
种属反应性	<b>与反应:</b> Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
表位	<b>ab32028</b> reacts with an epitope located in the C terminal region of mTOR.
阳性对照	IHC-P: Human breast carcinoma, mouse testis and rat testis tissues. IP: Rat brain tissue lysate and HeLa whole cell lysate ( <b>ab150035</b> ). IHC-Fr: Human heart tissue sections.
常规说明	<p>ab218525 is the carrier-free version of <b>ab32028</b>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <b><a href="#">see here</a></b>.</p>

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	Y391
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise<sup>™</sup>承诺保证使用ab218525于以下的经测试应用**

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		1/200.
WB		Use at an assay dependent concentration. Detects a band of approximately 250 kDa (predicted molecular weight: 289 kDa).
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <a href="#">IHC antigen retrieval protocols</a> .

**应用说明**      Is unsuitable for Flow Cyt or ICC/IF.

## 靶标

**功能**      Kinase subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. Growth factor-stimulated mTORC1 activation involves AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potentially activates the protein kinase activity of mTORC1. Amino-acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates

S6K1 at 'Thr-421', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Phosphorylates MAF1 leading to attenuation of its RNA polymerase III-repressive function. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'.

#### 组织特异性

Expressed in numerous tissues, with highest levels in testis.

#### 序列相似性

Belongs to the PI3/PI4-kinase family.

Contains 1 FAT domain.

Contains 1 FATC domain.

Contains 7 HEAT repeats.

Contains 1 PI3K/PI4K domain.


#### 翻译后修饰

Autophosphorylated; when part of mTORC1 or mTORC2.

#### 细胞定位

Endoplasmic reticulum membrane. Golgi apparatus membrane. Mitochondrion outer membrane. Lysosome. Cytoplasm. Nucleus > PML body. Shuttles between cytoplasm and nucleus. Accumulates in the nucleus in response to hypoxia (By similarity). Targeting to lysosomes depends on amino acid availability and RRAGA and RRAGB.

#### 图片



**All lanes :** Anti-mTOR antibody [Y391] ([ab32028](#)) at 1/1000 dilution

**Lane 1 :** Wild-type A549 cell lysate

**Lane 2 :** MTOR [homo] CRISPR-Cas9 edited A549 cell lysate

**Lane 3 :** HepG2 cell lysate

**Lane 4 :** HEK-293 cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 289 kDa

**Observed band size:** 250 kDa

False colour image of Western blot: Anti-mTOR antibody [Y391] staining at 1/1000 dilution, shown in green; Mouse anti-CANX [CANX/1543] ([ab238078](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab32028](#) was shown to bind


specifically to mTOR. A band was observed at 250 kDa in wild-type A549 cell lysates with no signal observed at this size in MTOR CRISPR-Cas9 edited cell line [ab283257](#). The band observed in the CRISPR-Cas9 edited lysate lane below 250 kDa is likely to represent a truncated form of mTOR. This has not been investigated further and the functional properties of the gene product have not been determined. To generate this image, wild-type and MTOR CRISPR-Cas9 edited A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween<sup>®</sup> 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-mTOR antibody [Y391] - BSA and Azide free ([ab218525](#))

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labelling mTOR with purified [ab32028](#) at a dilution of 1/400. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab32028](#)).




Immunohistochemistry (Frozen sections) - Anti-mTOR antibody [Y391] - BSA and Azide free (ab218525)

This data was developed using the same antibody clone in a different buffer formulation (**ab32028**).

IHC image of mTOR staining in a section of frozen normal human heart performed on a Leica BOND™ system using the standard protocol. The section was fixed in 10% paraformaldehyde (10 min) prior to staining. The section was incubated with **ab32028**, 1/200 dilution, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunoprecipitation - Anti-mTOR antibody [Y391] - BSA and Azide free (ab218525)

**ab32028** (purified) at a dilution of 1/100 immunoprecipitating mTOR in HeLa whole cell lysate.

Lane 1 (input): HeLa whole cell lysate (10µg)

Lane 2 (+): **ab32028** + HeLa whole cell lysate.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab32028** in HeLa whole cell lysate.

For western blotting, **ab131366** VeriBlot for IP (HRP) was used for detection at 1/1000 dilution.

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32028**).



Immunoprecipitation - Anti-mTOR antibody [Y391] - BSA and Azide free (ab218525)

**ab32028** (purified) at a dilution of 1/100 immunoprecipitating mTOR in rat brain tissue lysate.

Lane 1 (input): Rat brain tissue lysate (10µg)

Lane 2 (+): **ab32028** + rat brain tissue lysate.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab32028** in rat brain tissue lysate.

For western blotting, **ab131366** VeriBlot for IP (HRP) was used for detection at 1/1000 dilution.

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32028**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-mTOR antibody [Y391] - BSA and Azide free (ab218525)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse testis tissue labelling mTOR with purified **ab32028** at a dilution of 1/400. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32028**).





Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat testis tissue labelling mTOR with purified **ab32028** at a dilution of 1/400. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32028**).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-mTOR antibody [Y391] - BSA and Azide free (ab218525)

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-mTOR antibody [Y391] - BSA and Azide free (ab218525)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors