

### Anti-MUC16 antibody [EPR1020(2)] ab110640

**重组** RabMAb

★★★★★ **3 Abreviews** **6 References** **14 图像**

#### 概述

产品名称	Anti-MUC16抗体[EPR1020(2)]
描述	兔单克隆抗体[EPR1020(2)] to MUC16
宿主	Rabbit
经测试应用	<b>适用于:</b> WB, IHC-P
种属反应性	<b>与反应:</b> Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human uterus cell lysate. IHC-P: Human serous carcinoma and mucinous carcinoma tissues. ICC/IF: SK-OV-3 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS, 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆

克隆编号	EPR1020(2)
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™** 承诺保证使用ab110640于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/10000. Predicted molecular weight: 1519 kDa.
IHC-P	★★★★★ (3)	1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. <b>For unpurified use at 1/100 - 1/250.</b>

靶标


功能	Thought to provide a protective, lubricating barrier against particles and infectious agents at mucosal surfaces.
组织特异性	Expressed in corneal and conjunctival epithelia (at protein level). Overexpressed in ovarian carcinomas and ovarian low malignant potential (LMP) tumors as compared to the expression in normal ovarian tissue and ovarian adenomas.
序列相似性	Contains 2 ANK repeats. Contains 56 SEA domains.
结构域	Composed of three domains, a Ser-, Thr-rich N-terminal domain, a repeated domain containing more than 60 partially conserved tandem repeats of 156 amino acids each (AAs 12061-21862) and a C-terminal transmembrane contain domain with a short cytoplasmic tail.
翻译后修饰	Heavily O-glycosylated; expresses both type 1 and type 2 core glycans. Heavily N-glycosylated; expresses primarily high mannose and complex bisecting type N-linked glycans. May be phosphorylated. Phosphorylation of the intracellular C-terminal domain may induce proteolytic cleavage and the liberation of the extracellular domain into the extracellular space. May contain numerous disulfide bridges. Association of several molecules of the secreted form may occur through interchain disulfide bridges providing an extraordinarily large gel-like matrix in the extracellular space or in the lumen of secretory ducts.
细胞定位	Cell membrane. Secreted > extracellular space. May be liberated into the extracellular space following the phosphorylation of the intracellular C-terminus which induces the proteolytic cleavage and liberation of the extracellular domain.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human ovarian serous papillary carcinoma tissue sections labeling MUC16 with Purified ab110640 at 1:1000 dilution (0.12 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

**All lanes :** Anti-MUC16 antibody [EPR1020(2)] (ab110640) at 1/20000 dilution (purified)

**Lane 1 :** Human uterus lysates

**Lane 2 :** OVCAR-3 (Human ovary adenocarcinoma epithelial cell) whole cell lysates


Lysates/proteins at 15 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 1519 kDa


Blocking and diluting buffer: 5% NFDM/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 at 1/100 dilution staining MUC16 in Human serous carcinoma by Immunohistochemistry, Paraffin-embedded tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

**All lanes :** Anti-MUC16 antibody [EPR1020(2)] (ab110640) at 1/10000 dilution (purified)

**Lane 1 :** MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysates

**Lane 2 :** SK-OV-3 (Human ovarian cancer epithelial cell) whole cell lysates

**Lane 3 :** OVCAR-3 (Human ovary adenocarcinoma epithelial cell) whole cell lysates

Lysates/proteins at 15 µg per lane.


### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 1519 kDa

Blocking and diluting buffer: 5% NFDM/TBST


According to PMID: 26918940 and PMID: 19747716, MCF7 and SK-OV-3 cells are MUC16 negative cell lines



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 at 1/100 dilution staining MUC16 in Human mucinous carcinoma by Immunohistochemistry, Paraffin-embedded tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing positive staining in Serous ovarian carcinoma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing positive staining in Normal uterus tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing negative staining in Cervical carcinoma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing negative staining in Colonic adenocarcinoma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing negative staining in Normal kidney tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing negative staining in Normal colon tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing negative staining in Normal breast tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MUC16 antibody [EPR1020(2)] (ab110640)

Unpurified ab110640 showing negative staining in Normal brain tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

#### Why choose a recombinant antibody?



Anti-MUC16 antibody [EPR1020(2)] (ab110640)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors