

Anti-Muscarinic Acetylcholine Receptor M4/CHRM4 antibody [18C7.2] ab77956

★★★★★ [2 Abreviews](#) [5 References](#) [1 图像](#)

概述

产品名称	Anti-Muscarinic Acetylcholine Receptor M4/CHRM4抗体[18C7.2]
描述	小鼠单克隆抗体[18C7.2] to Muscarinic Acetylcholine Receptor M4/CHRM4
宿主	Mouse
特异性	ab77956 reacts with Muscarinic Acetylcholine Receptor M4/CHRM4. There is no reactivity with the other subtypes.
经测试应用	适用于: IHC-Fr, IP, Flow Cyt, WB
种属反应性	与反应: Mouse, Rat, Human, Monkey
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Transfected CHO cell lines
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.1% Sodium azide Constituents: 1.45% Sodium chloride, 0.0536% PBS
纯度	Protein A purified
纯化说明	Purified from TCS.
克隆	单克隆
克隆编号	18C7.2
同种型	IgG1

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab77956于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration.
WB	★★★★★ (2)	Use at an assay dependent concentration.

靶标

图片

Flow Cytometry - Anti-Muscarinic Acetylcholine
Receptor M4/CHRM4 antibody [18C7.2] (ab77956)

Overlay histogram showing SH-SY5Y cells stained with ab77956 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab77956, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.

This image was generated using the ascites version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors