abcam

Product datasheet

Anti-Myc tag antibody [9E10] ab206486



★★★★★ 4 Abreviews 15 References 4 图像

概述

产**品名称** Anti-Myc tag抗体[9E10]

描述 大鼠单克隆抗体[9E10] to Myc tag

宿主 Rat

特异性 This antibody is specific for Myc tagged proteins. The Myc tag epitope (EQKLISEEDL) is located

at the dimerization site of c-myc and therefore this antibody does not perform well at recognizing

endogenous c-myc.

经测试应用 适用于: IHC-Fr, IHC-P, ICC/IF, IP, WB

免疫原 Synthetic peptide corresponding to Human Myc tag aa 400 to the C-terminus (C terminal).

Database link: P01106

Run BLAST with
Run BLAST with

表位 Epitope located at aa 410-419; EQKLISEEDL

常规说明 ab206486 is a recombinant antibody that comprises the variable regions from the 9E10

hybridoma within a rat lgG1 scaffold.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

存储溶液 Preservative: 0.02% Proclin 300

Constituent: 99% PBS

纯**度** Protein A purified

 克隆
 单克隆

 克隆编号
 9E10

 同种型
 IgG1

 轻链类型
 kappa

应用

1

The Abpromise guarantee

Abpromise™承诺保证使用ab206486于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
ICC/IF	*****(1)	Use at an assay dependent concentration.
IP	★★★★ (1)	Use at an assay dependent concentration.
WB	★★★★☆ (1)	Use at an assay dependent concentration. Predicted molecular weight: 48 kDa.

靶标


相关性

Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells.

细胞定位

Nuclear

图片



Western blot - Anti-Myc tag antibody [9E10] (ab206486)

All lanes: Anti-Myc tag antibody [9E10] (ab32) at 1/200 dilution

Lane 1: Wild-type HEK-293T cell lysate

Lane 2: MYC CRISPR-Cas9 edited HEK-293T cell lysate

Lane 3 : Jurkat cell lysate


Lane 4 : SH-SY5Y cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 48 kDa

False colour image of Western blot: Anti-Myc tag antibody [9E10] staining at 1/200 dilution, shown in green; Rabbit Anti-GAPDH antibody [EPR16891] (ab181602) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab32 was shown to bind specifically to Myc tag. A band was observed at 57 kDa in wild-type HEK-293T cell lysates with no signal observed at this size in MYC CRISPR-Cas9 edited cell line ab256500 (CRISPR-Cas9 edited cell lysate ab263850). The band observed in the CRISPR-Cas9 edited lysate lane below 57 kDa is likely to represent a truncated form of Myc tag. This has not been investigated further and the functional properties of the gene product have not been determined. To generate this image, wild-type and MYC CRISPR-Cas9 edited HEK-293T cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed (ab216772) and Goat anti-Rabbit lgG H&L (IRDye® 680RD) preabsorbed (ab216777) at 1/20000 dilution.



Western blot - Anti-Myc tag antibody [9E10] (ab206486)



All lanes: Anti-Myc tag antibody [9E10] (ab206486) at 2 µg/ml


Lane 1: HEK-293 spiked with 5 ng multitag protein

Lane 2: HEK-293 unspiked with 5 ng multitag protein

Lane 3: 25 ng purified multitag protein

Lysates/proteins at 35 µg per lane.

Predicted band size: 48 kDa



Western blot - Anti-Myc tag antibody [9E10] (ab206486)


All lanes: Anti-Myc tag antibody [9E10] (ab206486) at 1 µg/ml

Lane 1: Multitag protein at 0.025 µg

Lane 2: Multitag protein at 0.01 µg

Lane 3: Multitag protein at 0.005 µg

Predicted band size: 48 kDa



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors