

Anti-NCR1 antibody [n1D9] ab14823

3 References **3 图像**

概述

产品名称	Anti-NCR1抗体[n1D9]
描述	小鼠单克隆抗体[n1D9] to NCR1
宿主	Mouse
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Human
免疫原	Recombinant full length protein, corresponding to amino acids 22-255 of Human NCR1
阳性对照	Flow Cyt: Human peripheral blood mononuclear cells or natural killer cells, PBMC cells, HeLa cells.
常规说明	<p>This product was changed from ascites to tissue culture supernatant on 24/01/2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: PBS, 10% Glycerol</p>
纯度	Tissue culture supernatant
纯化说明	Purified from TCS.
克隆	单克隆
克隆编号	n1D9

骨髓瘤	Sp2/0
同种型	IgG1
轻链类型	kappa

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab14823于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use at an assay dependent concentration. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.


靶标

功能	Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis.
组织特异性	Selectively expressed by both resting and activated NK cells.
序列相似性	Belongs to the natural cytotoxicity receptor (NCR) family. Contains 2 Ig-like (immunoglobulin-like) domains.
翻译后修饰	N-glycosylated. O-glycosylated.
细胞定位	Cell membrane.

图片




Flow cytometry analysis of NKp46 in HeLa cells. The cell was stained with ab14823 at 2-5µg for 1x10⁶ cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (blue), cells without incubation with primary and secondary antibody was used as the negative control (black).



Flow Cytometry - Anti-NCR1 antibody [n1D9]
(ab14823)

Flow cytometry analysis of NKp46 in PBMC cells. The cell was stained with ab14823 at 2-5 μ g for 1x10⁶ cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (blue), cells without incubation with primary and secondary antibody was used as the negative control (black).



Flow Cytometry - Anti-NCR1 antibody [n1D9]
(ab14823)

ab14823 staining human PBMC by flow cytometry.

This image was produced using the ascites version of this antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors