




Anti-nNOS (neuronal) (phospho S1417) antibody ab5583

★★★★★ **6 Abreviews** **20 References** **2 图像**


概述

产品名称	Anti-nNOS (neuronal) (phospho S1417)抗体
描述	兔多克隆抗体 to nNOS (neuronal) (phospho S1417)
宿主	Rabbit
经测试应用	适用于: IHC-Fr, ICC, WB, IP, ICC/IF, IHC-FoFr
种属反应性	与反应: Mouse, Rat, Rabbit, Human 预测可用于: Xenopus laevis, Zebrafish 
免疫原	Synthetic peptide corresponding to Human nNOS (neuronal) aa 1411-1425 (phospho S1417). Sequence: CNRLRSE(pS)IAFIEESK Database link: P29475 (Peptide available as ab41773)  Run BLAST with  Run BLAST with
阳性对照	WB: nocozadole-treated SH-SY5Y cells; ICC: SH-SY5Y cells
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 99% PBS
纯度	Immunogen affinity purified
克隆	多克隆

同种型	IgG	
应用		
The Abpromise guarantee Abpromise™ 承诺保证使用ab5583于以下的经测试应用		
“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。		
应用	Ab评论	说明
IHC-Fr	★★★★★ (2)	Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.
WB	★★★★★ (4)	1/500 - 1/5000. Predicted molecular weight: 160 kDa.
IP		Use at an assay dependent concentration.
ICC/IF		1/10 - 1/100.
IHC-FoFr		Use at an assay dependent concentration. PubMed: 20881244
靶标		
功能	Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In the brain and peripheral nervous system, NO displays many properties of a neurotransmitter. Probably has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such SRR.	
组织特异性	Isoform 1 is ubiquitously expressed: detected in skeletal muscle and brain, also in testis, lung and kidney, and at low levels in heart, adrenal gland and retina. Not detected in the platelets. Isoform 3 is expressed only in testis. Isoform 4 is detected in testis, skeletal muscle, lung, and kidney, at low levels in the brain, but not in the heart and adrenal gland.	
序列相似性	Belongs to the NOS family. Contains 1 FAD-binding FR-type domain. Contains 1 flavodoxin-like domain. Contains 1 PDZ (DHR) domain.	
结构域	The PDZ domain in the N-terminal part of the neuronal isoform participates in protein-protein interaction, and is responsible for targeting nNos to synaptic membranes in muscles. Mediates interaction with VAC14.	
翻译后修饰	Ubiquitinated; mediated by STUB1/CHIP in the presence of Hsp70 and Hsp40 (in vitro).	
细胞定位	Cell membrane > sarcolemma. Cell projection > dendritic spine. In skeletal muscle, it is localized beneath the sarcolemma of fast-twitch muscle fiber by associating with the dystrophin glycoprotein complex. In neurons, enriched in dendritic spines.	
图片		



Western blot - Anti-nNOS (neuronal) (phospho S1417) antibody (ab5583)

All lanes : Anti-nNOS (neuronal) (phospho S1417) antibody (ab5583) at 1/1000 dilution

Lane 1 : untreated SH-SY5Y cells (Human neuroblastoma cell line from bone marrow)

Lane 2 : SH-SY5Y cells (Human neuroblastoma cell line from bone marrow) treated with 100 ng/mL nocodazole


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : HRP-conjugated secondary antibody

Developed using the ECL technique.

Predicted band size: 160 kDa



Immunocytochemistry/ Immunofluorescence - Anti-nNOS (neuronal) (phospho S1417) antibody (ab5583)

Immunocytochemistry/Immunofluorescence analysis of nNOS (neuronal) (phospho S1417) (green) showing staining in the cytoplasm and nucleus of SH-SY5Y cells treated with 100 ng/ml nocodazole (left) and untreated SH-SY5Y cells (right). Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were incubated with ab5583 in 3% BSA-PBS at a dilution of 1:20 and incubated overnight at 4°C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight-conjugated secondary antibody in PBS at room temperature in the dark. F-actin (red) was stained with a fluorescent red phalloidin and nuclei (blue) were stained with Hoechst or DAPI. Images were taken at a magnification of 60x.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors