abcam

Product datasheet

PE Anti-Estrogen Receptor alpha antibody [E115] ab209288



重组 RabMAb

★★★★★ 3 Abreviews 2 References 5 图像

概述

产品名称 PE Anti-Estrogen Receptor alpha抗体[E115]

描述 PE兔单克隆抗体[E115] to Estrogen Receptor alpha

宿主 Rabbit

偶联物 PE. Ex: 488nm. Em: 575nm

特异性 Expression levels of ER alpha protein vary with sample type.

经测试应用 适用于: Flow Cyt (Intra), ICC/IF

不适用于: IHC-P

种属反应性 与反应: Mouse, Rat, Human

预测可用于: Dog 🕰

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 Flow Cyt (Intra): T47D, 4T1 and GH3 cells. ICC/IF: T47D cells.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式

Shipped at 4°C. Upon delivery aliquot. Store at +4°C. Do Not Freeze. Store In the Dark. 存放说明

存储溶液

Preservative: 0.02% Sodium azide Constituents: PBS, 1% BSA

纯度 Protein A purified

克隆 单克隆

克隆编号 E115

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab209288于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/50 - 1/2500.
ICC/IF		1/500. This product gave a positive signal in T47D cells fixed with 4% formaldehyde (10 min)

应用说明 Is unsuitable for IHC-P.

靶标

功能 Nuclear hormone receptor. The steroid hormones and their receptors are involved in the

regulation of eukaryotic gene expression and affect cellular proliferation and differentiation in

target tissues. Can activate the transcriptional activity of TFF1.

序列相似性 Belongs to the nuclear hormone receptor family. NR3 subfamily.

Contains 1 nuclear receptor DNA-binding domain.

结构域 Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-

terminal ligand-binding domain.

翻译后修饰 Phosphorylated by cyclin A/CDK2. Phosphorylation probably enhances transcriptional activity.

Glycosylated; contains N-acetylglucosamine, probably O-linked.

Ubiquitinated. Deubiquitinated by OTUB1.


Dimethylated by PRMT1 at Arg-260. The methylation may favor cytoplasmic localization.

Palmitoylated (isoform 3). Not biotinylated (isoform 3).

细胞定位 Nucleus. Cytoplasm. Cell membrane. A minor fraction is associated with the inner membrane and

Nucleus. Cytoplasm. Cell membrane. Associated with the inner membrane via palmitoylation.


图片



Immunocytochemistry/ Immunofluorescence - PE
Anti-Estrogen Receptor alpha antibody [E115]
(ab209288)

ab209288 staining Estrogen Receptor alpha in T47D cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab209288 at 1/500 dilution (Pseudocolored in green) and ab195884, Rat monoclonal to Tubulin (Alexa Fluor® 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Flow Cytometry (Intracellular) - PE Anti-Estrogen Receptor alpha antibody [E115] (ab209288)

Overlay histogram showing T47D cells stained with ab209288 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilized with 90% methanol for 30 min at -20°C. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab209288, 1/2500 dilution) for 30 min at 22°C.


Isotype control antibody (black line) was rabbit IgG (monoclonal) Phycoerythrin (<u>ab209478</u>) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 50 mW Yellow/Green laser (561nm) and 586/15 bandpass filter.




Flow Cytometry (Intracellular) - PE Anti-Estrogen Receptor alpha antibody [E115] (ab209288)

Intracellular Flow Cytometry analysis of 4T1 (mouse mammary gland carcinoma epithelial cell) cells labeling Estrogen Receptor alpha with ab209288 at 1/50 dilution (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. Isotype control - Rabbit IgG monoclonal [EPR25A, ab209478] (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Flow Cytometry (Intracellular) - PE Anti-Estrogen Receptor alpha antibody [E115] (ab209288)

Intracellular Flow Cytometry analysis of GH3 (rat pituitary epithelial) cells labeling Estrogen Receptor alpha with ab209288 at 1/50 dilution (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. Isotype control - Rabbit IgG monoclonal [EPR25A, <u>ab209478</u>] (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors