

PE Anti-TRAIL antibody [N2B2] ab95532

3 References **1 图像**

概述

产品名称	PE Anti-TRAIL抗体[N2B2]
描述	PE大鼠单克隆抗体[N2B2] to TRAIL
宿主	Rat
偶联物	PE. Ex: 488nm, Em: 575nm
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Mouse
免疫原	Tissue, cells or virus corresponding to Mouse TRAIL. Mouse TRAIL-transfected 2PK-3 cells. Database link: P50592
阳性对照	Flow Cyt: Mouse TRAIL transfected cells.
常规说明	<p>ab95532 inhibits IL-2 or IL-15 activated NK cell cytotoxicity against mouse fibrosarcoma L929 target cells.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	pH: 7.20 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, PBS
纯度	Protein G purified
Primary antibody说明	ab95532 inhibits IL-2 or IL-15 activated NK cell cytotoxicity against mouse fibrosarcoma L929 target cells.

克隆	单克隆
克隆编号	N2B2
同种型	IgG2a
轻链类型	kappa

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab95532于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use 0.06-0.125µg for 10 ⁵⁻⁸ cells. Staining the cell sample in a final volume of 100µl is recommended.

靶标

功能	Cytokine that binds to TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and possibly also to TNFRSF11B/OPG. Induces apoptosis. Its activity may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4 and TNFRSF11B/OPG that cannot induce apoptosis.
组织特异性	Widespread; most predominant in spleen, lung and prostate.
序列相似性	Belongs to the tumor necrosis factor family.
细胞定位	Membrane.

图片



Flow cytometry staining of TRAIL-transfected cells with 0.06 µg of Rat IgG2a Isotype Control PE (pink histogram) or 0.06 µg of ab95532 (green histogram). Total viable cells were used for analysis.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors